

42353 D B2C2

No. 14

① Europäisches Patentamt
European Patent Office
Office européen des brevets

⑪ Publication number:

0 029 557
A2

⑫ EUROPEAN PATENT APPLICATION

⑬ Application number: 80107074.9

⑭ Date of filing: 15.11.80

⑮ Int. Cl. 3: C 07 D 501/36
C 07 D 501/34, C 07 D 501/30
C 07 D 501/32
//C07D209/48, C07D277/20

⑯ Priority: 19.11.79 GB 7939985
08.02.80 GB 8004335
21.04.80 GB 8012991

⑰ Date of publication of application:
03.06.81 Bulletin 81/22

⑱ Designated Contracting States:
AT BE CH DE FR GB IT LI NL SE

⑲ Applicant: Fujisawa Pharmaceutical Co. Ltd.
3, Doshomachi 4-chome Higashi-ku
Osaka-shi, Osaka 541(JP)

⑳ Inventor: Takaya, Takeo
No. 1-5-87, Suimeidai
Kawanishi(JP)

㉑ Inventor: Takasugi, Hisashi
No. 2-12-7, Kohama-nishi
Suminoe Osaka(JP)

㉒ Inventor: Masugi, Takashi
No. 3-10-11, Hachizuka
Ikeda(JP)


㉓ Inventor: Yamanaka, Hideaki
No. 2-77-10, Nakanoshima
Kuzuhara Hirakata(JP)

㉔ Inventor: Kawabata, Kohji
No. 1-43-2, Orion-cho
Sumiyoshi Osaka(JP)


㉕ Representative: Türk, Dietmar, Dr. rer. nat. et al.
Redies, Redies, Türk & Gille Bruckner Strasse 20
D-4000 Düsseldorf 13(DE)

① 7-Acylamino-3-substituted cephalosporanic acid derivatives, processes for their preparation, pharmaceutical compositions containing them; their starting compounds and their preparation.

② 7-Acylamino-3-substituted cephalosporanic acid derivatives of the formula:



in which R¹ is a group of the formula:



wherein R⁴ is lower alkyl and
R⁵ is amino or a protected amino group,
R² is lower alkoxymethyl, lower alkyl-
thiomethyl or lower alkenylthiomethyl,
R³ is carboxy or a protected carboxy group, and
A is lower alkylene which may have certain substituents

and pharmaceutically acceptable salt thereof and processes for their preparation and also a pharmaceutically composition comprising, as an effective ingredient, the above compound in association with pharmaceutically acceptable carriers. The invention also relates to the starting compounds

7-ACYLAMINO - 3-SUBST.
CEPHALOSPORIN DERIVS. + USEFUL
AS ANTIBACTERIALS IN HUMAN
AND VETERINARY MEDICINE


EP 0 029 557 A2

769

42353 D

Croydon Printing Company Ltd.

453



and their preparation.

TITLE MODIFIED
See front page

-1-

7-ACYLAMINO-3-SUBSTITUTED CEPHALOSPORANIC ACID
DERIVATIVES, PROCESSES FOR THEIR PREPARATION,
PHARMACEUTICAL COMPOSITIONS CONTAINING THEM
AND THEIR STARTING COMPOUNDS

The present invention relates to novel 7-acylamino-3-substituted cephalosporanic acid derivatives and pharmaceutically acceptable salts thereof.

More particularly, it relates to novel 7-acylamino-3-substituted cephalosporanic acid derivatives and pharmaceutically acceptable salts thereof, which have antimicrobial activity, to processes for the preparation thereof, to a pharmaceutical composition comprising the same, and to a method of using the same therapeutically in the treatment of infectious diseases in human being and animals.

Accordingly, one object of the present invention is to provide novel 7-acylamino-3-substituted cephalosporanic acid derivatives and pharmaceutically acceptable

15

salts thereof, which are highly active against a number of pathogenic microorganisms and are useful as antimicrobial agents, especially oral administration.


Another object of the present invention is to

provide processes for the preparation of novel 7-acylamino-3-substituted cephalosporanic acid derivatives and salts thereof.


A further object of the present invention is to provide a pharmaceutical composition comprising, as active ingredients, said 7-acylamino-3-substituted cephalosporanic acid derivatives and pharmaceutically acceptable salts thereof.

Still further object of the present invention is to provide a method of using said 7-acylamino-3-substituted cephalosporanic acid derivatives and pharmaceutically acceptable salts thereof in the treatment of infectious diseases by pathogenic microorganisms in human being and animals.

The object 7-acylamino-3-substituted cephalosporanic acid derivatives are novel and can be represented by the following general formula:



in which R^1 is a group of the formula:



wherein R^4 is lower alkyl and

R^5 is amino or a protected amino group,

R^2 is lower alkoxyethyl, lower alkylthiomethyl or lower alkenylthiomethyl,


R^3 is carboxy or a protected carboxy group, and

A is lower alkylene which may have a substituent selected from the groups consisting of amino, a protected amino group, hydroxy, oxo and a group of the formula:


$\text{-N}^+\text{OR}^6$, wherein R^6 is hydrogen, lower alkenyl, lower alkynyl, lower alkyl or lower alkyl substituted by one or more substituent(s) selected from carboxy, a protected carboxy group, amino, a protected amino group, and heterocyclic group.

In the object compounds (I) and the corresponding starting compounds (II) to (IV) in Processes 1 and 7 mentioned below, it is to be understood that there may be one or more stereoisomeric pair(s) such as optical and geometrical isomers due to asymmetric carbon atom and double bond in those molecules and such isomers are also included within the scope of the present invention.

With regard to geometrical isomers in the object compounds and the starting compounds, it is to be noted that, for example, the object compounds, wherein A means a group of the formula: $\text{-C=N}^+\text{OR}^6$, include syn isomer, anti isomer and a mixture thereof, and the syn isomer means one geometrical isomer having the partial structure represented by the following formula:



and the anti isomer means the other geometrical isomer having the partial structure represented by the following formula:



Regarding the other object and starting compounds as mentioned above, the syn isomer and the anti isomer

42353D

772

456

773

42353D

457

can also be referred to the same geometrical isomers as illustrated for the compounds (I).


Suitable pharmaceutically acceptable salts of the object compounds (I) are conventional non-toxic salts and may include a salt with a base or an acid addition salt such as a salt with an inorganic base, for example, an alkali metal salt (e.g. sodium salt, potassium salt, etc.), an alkaline earth metal salt (e.g. calcium salt, magnesium salt, etc.), an ammonium salt; a salt with an organic base, for example, an organic amine salt (e.g. triethylamine salt, pyridine salt, picoline salt, ethanalamine salt, triethanolamine salt, dicyclohexylamine salt, N,N'-dibenzylethylenediamine salt, etc.) etc.; an inorganic acid addition salt (e.g. hydrochloride, hydrobromide, sulfate, phosphate, etc.); an organic carboxylic or sulfonic acid addition salt (e.g. formate, acetate, trifluoroacetate, maleate, tartrate, methanesulfonate, benzenesulfonate, p-toluenesulfonate, etc.); a salt with a basic or acidic amino acid (e.g. arginine, aspartic acid, glutamic acid, etc.); an intermolecular or intramolecular quaternary salt, and the like.

The said intermolecular quaternary salt can be formed in case that the heterocyclic group in R⁶ in the compounds (I) contains nitrogen atom(s) (e.g. pyridyl, etc.), and suitable intermolecular quaternary salt may include 1-lower alkylpyridinium lower alkylsulfate (e.g. 1-methylpyridinium methylsulfate, 1-ethylpyridinium ethylsulfate, etc.), 1-lower alkylpyridinium halide (e.g. 1-methylpyridinium iodide, etc.) and the like. The said intramolecular salt can be formed in case that heterocyclic group in R⁶ in the compounds (I) contains nitrogen atom(s) (e.g. pyridyl etc.) and R³ is carboxy, and suitable intramolecular salt may include 1-lower alkylpyridinium carboxylate (e.g. 1-methylpyridinium carboxylate, 1-ethylpyridinium

carboxylate, 1-propylpyridinium carboxylate, 1-isopropylpyridinium carboxylate, 1-butylpyridinium carboxylate, etc.); and the like.

According to the present invention, the object compounds (I) and the pharmaceutically acceptable salts thereof can be prepared by the processes as illustrated by the following reaction schemes.

10 (1) Process 1:



(II)

or its reactive derivative at the amino group or a salt thereof

(I)


or a salt thereof

(continued to the next page)


0029557

-7-


(2) Process 2:




(3) Process 3:




(4) Process 4:




(5) Process 5:




(6) Process 6:



(7) Process 7:



(8) Process 8:



(continued to the next page)

0029557

-6-

776 423530


460

423530

777

461

(9) Process 9:




(I-m)

or a salt thereof

(I-n)

or a salt thereof

10 (10) Process 10:




(I-o)

or a salt thereof

(I-p)

or a salt thereof

20 (11) Process 11:




(I-l)

or a salt thereof

(I-k)

or a salt thereof

(12) Process 12:




(I-q)

35 or a salt thereof

(I-r)

or a salt thereof

(13) Process 13:



(I-r)

or a salt thereof



(I-s)

or a salt thereof

10 in which R¹, R², R³, R⁵

and A are each as defined

above,

R¹_a is a group of the formula:wherein R⁸ is a protected amino group,R¹_b is a group of the formula:R³_a is a protected carboxy group,R³_b is lower alkoxy-carbonyl substituted by protected amino and protected carboxy groups,R³_c is lower alkoxy-carbonyl substituted by amino and carboxy,R⁷ is lower alkyl,A¹ is lower alkylene having a protected amino group,A² is lower alkylene having an amino group,A³ is lower alkylene having an oxo group,A⁴ is lower alkylene having a hydroxy group,A⁵ is lower alkylene which may have

a group of the formula $-N^{\circ}OR^6$, wherein R^6 is as defined above,

A^6 is lower alkylene having a group of the formula: $-N^{\circ}OR^6_a$, wherein R^6_a is lower alkyl substituted by a protected carboxy group,


A^7 is lower alkylene having a group of the formula: $-N^{\circ}OR^6_b$, wherein R^6_b is lower alkyl substituted by carboxy,

A^8 is lower alkylene having a group of the formula: $-N^{\circ}OR^6_c$, wherein R^6_c is lower alkoxy-carbonyl(lower)alkyl substituted by protected amino and protected carboxy groups or lower alkyl substituted by protected amino and protected carboxy groups,

A^9 is lower alkylene having a group of the formula: $-N^{\circ}OR^6_d$, wherein R^6_d is lower alkoxy-carbonyl(lower)alkyl substituted by amino and carboxy or lower alkyl substituted by amino and carboxy,


A^{10} is lower alkylene having a group of the formula: $-N^{\circ}OR^6_e$, wherein R^6_e is lower alkyl substituted by a group of the formula:

A^{11} is lower alkylene having a group of the formula: $-N^{\circ}OR^6_f$, wherein R^6_f is lower alkyl substituted by a group of the formula:



wherein R^7 is as defined above,

A^{12} is lower alkylene having a group of the formula: $-N^{\circ}OR^6_g$, wherein R^6_g is lower alkyl substituted by a cation of the formula:




wherein R^7 is as defined above, and

X^1 is halogen.

Some of the starting compounds (II), (III) and (IV) used in Processes 1 and 7 are new and can be

represented by the following formula:



in which RA is amino or a protected amino group, RB' is lower alkenyl, X is $-S-$ or $-SO-$, and R^3 is as defined above, and a salt thereof.


Suitable salts of the above starting compound are the same as those exemplified for the object compounds (I). The starting compound thus formulated and other starting compounds can be

(continued to the next page)

prepared, for example, from the known compounds by the methods in the following reaction schemes, and others can be prepared in a similar manner thereto or in a conventional manner.

5

(A) Process A: Preparation of some of the starting compounds (II)



15


(II-a)

or a salt thereof

(II-b)

or a salt thereof

20

Reduction
(Process A-2)

25

(II-d)

or a salt thereof


(II-c)

or a salt thereof

30

35

(B) Process B: Preparation of some of the starting compounds (III)



782


42353D

783

42353D

466

467

(C) Process C: Preparation of some of the starting compounds (IV)

10


(IV-a)

or its reactive derivative at the amino group or a salt thereof

15

(continued to the next page)

20

(D) Process b: Preparation of some of the compounds (II)

10

(II-e)

or a reactive derivative at the carboxy group or a salt thereof

15

Removal of the amino-protective group for R^a (Process D-2)

20

(II-f)

or a salt thereof



25

(E) Process E: Preparation of some of the compound (III) in Process 1

30

(III-h)

35





25

(E) Process E: Preparation of some of the compound (III) in Process 1

30

(III-h)

35



42353D

In the above and subsequent description of the present specification, suitable examples and illustration of the various definitions to be included within the scope thereof are explained in detail as follows.

The term "lower" in the present specification is intended to mean a group having 1 to 6 carbon atoms, unless otherwise indicated.

Suitable "lower alkyl" group may include straight or branched one such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, isopentyl, neopentyl, hexyl and the like, in which the preferred one is C_1 - C_3 alkyl.

Suitable "lower alkenyl" group may include straight or branched one such as vinyl, 1-propenyl, allyl, 1-(or 2- or 3-)butenyl, 1-(or 2- or 3- or 4-)pentenyl, 1-(or 2- or 3- or 4- or 5-)hexenyl, 2-methyl-2-propenyl, and the like, in which the preferred one is C_2 - C_5 alkenyl.

Suitable "lower alkynyl" group may include straight or branched one such as propargyl, 2-(or 3-)butynyl, 2-(or 3- or 4-)pentynyl, 2-(or 3- or 4- or 5-)hexynyl, and the like, in which the preferred one is C_2 - C_5 alkynyl.

(continued to the next page)

423530

423530

Suitable "lower alkoxyethyl" group may include a methyl group substituted by straight or branched lower alkoxy such as methoxymethyl, ethoxymethyl, propoxymethyl, isopropoxymethyl, pentyloxymethyl, hexyloxymethyl, and the like, in which the preferred one is C_1 - C_3 alkoxymethyl.

Suitable "lower alkylthiomethyl" group may include a methyl group substituted by straight or branched lower alkylthio such as methylthiomethyl, ethylthiomethyl, propylthiomethyl, isobutylthiomethyl, pentylthiomethyl, hexylthiomethyl, and the like, in which the preferred one is C_1 - C_3 alkylthiomethyl.

Suitable "lower alkenylthiomethyl" group may include a methyl group substituted by straight or branched lower alkenylthio such as vinylthiomethyl, 1-propenylthiomethyl, 1-allylthiomethyl, 2-methyl-2-propenylthiomethyl, 1-(or 2- or 3-)butenylthiomethyl, 1-(or 2- or 3- or 4-)pentenylthiomethyl, 1-(or 2- or 3- or 4-)hexenylthiomethyl, and the like, in which the preferred one is C_2 - C_5 alkenylthiomethyl.

Suitable "protected amino group" may include an

amino group substituted by a conventional amino-protective group which is used in penicillin and cephalosporin compounds, for example, acyl as mentioned below, mono- (or di or tri)phenyl(lower)alkyl (e.g. benzyl, benzhydryl, trityl, etc.), lower alkoxy carbonyl(lower)-alkylidene or its enamine tautomer (e.g. 1-methoxycarbonyl-1-propen-2-yl, etc.), di(lower)alkylamino-methylene (e.g. dimethylaminomethylene, etc.), etc.

Suitable "acyl" may include an aliphatic acyl, an aromatic acyl, a heterocyclic acyl and an aliphatic acyl substituted with aromatic or heterocyclic group(s).

The aliphatic acyl may include saturated or

unsaturated, acyclic or cyclic ones, such as lower alkanoyl (e.g. formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl,

etc.), lower alkanesulfonyl (e.g. mesyl, ethanesulfonyl, propanesulfonyl, etc.), lower alkoxycarbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, tert-butoxycarbonyl, etc.), lower alkenoyl (e.g. acryloyl, methacryloyl, crotonoyl, etc.), (C₃-C₇)-cycloalkanesulfonyl (e.g. cyclohexanesulfonyl, etc.), amidino, and the like.

The aromatic acyl may include aroyl (e.g. benzoyl, toluoyl, xyloyl, etc.), arenesulfonyl (e.g. benzene-sulfonyl, tosyl, etc.), and the like.

The heterocyclic acyl may include heterocycle-carbonyl (e.g. furoyl, thenoyl, nicotinoyl, isonicotinoyl, thiazolylcarbonyl, thiadiazolylcarbonyl, tetrazolylcarbonyl, etc.), and the like.

The aliphatic acyl substituted with aromatic group(s) may include phenyl(lower)alkanoyl (e.g. phenylacetyl, phenylpropionyl, phenylhexanoyl, etc.), phenyl(lower)-alkoxycarbonyl (e.g. benzyloxycarbonyl, phenethyloxy-carbonyl, etc.), phenoxy(lower)alkanoyl (e.g. phenoxy-acetyl, phenoxypropionyl, etc.), and the like.

The aliphatic acyl substituted with heterocyclic group(s) may include thienylacetyl, imidazolylacetyl, furylacetyl, tetrazolylacetyl, thiazolylacetyl, thiadiazolylacetyl, thienylpropionyl, thiadiazolyl-propionyl, and the like.

These acyl groups may be further substituted with one or more suitable substituents such as lower alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, pentyl, hexyl, etc.), halogen (e.g. chlorine, bromine, iodine, fluorine), lower alkoxy (e.g. methoxy, ethoxy, propoxy, isopropoxy, butoxy, pentyloxy, hexyloxy, etc.), lower alkylthio (e.g. methylthio, ethylthio, propylthio, isopropylthio, butylthio, pentylthio, hexylthio, etc.), nitro and the like, and preferable acyl having such 35 substituent(s) may be mono (or di or tri)halo(lower)-

42353D

alkanoyl (e.g. chloroacetyl, bromoacetyl, dichloroacetyl, trifluoroacetyl, etc.), mono (or di or tri)halo(lower)-alkoxycarbonyl (e.g. chloromethoxycarbonyl, dichloromethoxycarbonyl, 2,2,2-tri-chloroethoxycarbonyl, etc.), nitro (or halo or lower alkoxy)phenyl(lower)alkoxy-carbonyl (e.g. nitrobenzyloxycarbonyl, chlorobenzyloxy-carbonyl, methoxybenzyloxycarbonyl, etc.), and the like.

Suitable "protected carboxy group" may include an esterified carboxy group which is conventionally used 10 in penicillin or cephalosporin compounds at their 3rd or 4th position thereof.

Suitable "ester moiety" in "esterified carboxy group" may include lower alkyl ester (e.g. methyl ester, ethyl ester, propyl ester, isopropyl ester, butyl ester, 15 isobutyl ester, t-butyl ester, pentyl ester, tert-pentyl ester, hexyl ester, etc.), lower alkenyl ester (e.g. vinyl ester, allyl ester, etc.), lower alkynyl ester (e.g. ethynyl ester, propynyl ester, etc.), lower alkoxy(lower)alkyl ester (e.g. methoxymethyl ester, 20 ethoxymethyl ester, isopropoxymethyl ester, 1-methoxyethyl ester, 1-ethoxyethyl ester, etc.), lower alkylthio(lower)alkyl ester (e.g. methylthiomethyl ester, ethylthiomethyl ester, ethylthioethyl ester

isopropylthiomethyl ester, etc.), amino- and carboxy-substituted lower alkyl ester (e.g. 2-amino-2-carboxyethyl ester, 3-amino-3-carboxypropyl ester, etc.), protected amino and protected carboxy substituted lower alkyl ester such as lower alkoxycarbonylamino and mono- (or di or tri)phenyl(lower)alkoxycarbonyl substituted lower 30 alkyl ester (e.g. 2-tert-butoxycarbonylamino-2-benzhydryloxycarbonyl ethyl, 3-tert-butoxycarbonylamino-3-benzhydryloxycarbonylpropyl, etc.), mono(or di or tri)halo(lower)alkyl ester (e.g. 2-iodoethyl ester, 2,2,2-trichloroethyl ester, etc.), lower alkanoyloxy(lower)-

35

42353D

alkyl ester (e.g. acetoxymethyl ester, propionyloxy-
methyl ester, butyryloxymethyl ester, isobutyryloxy-
methyl ester, valeryloxymethyl ester, pivaloyloxymethyl-
ester, hexanoyloxymethyl ester, 2-acetoxyethyl ester,
5 2-propionyloxyethyl ester, 1-acetoxypropyl ester,
etc.), lower alkanesulfonyl(lower)alkyl ester (e.g.
mesylmethyl ester, 2-mesyloethyl ester, etc.), mono(or
di or tri)phenyl(lower)alkyl ester which may have one
or more suitable substituent(s) (e.g. benzyl ester, 4-
10 methoxybenzyl ester, 4-nitrobenzyl ester, phenethyl
ester, benzhydryl ester, trityl ester, bis(methoxyphenyl)-
methyl ester, 3,4-dimethoxybenzyl ester, 4-hydroxy-3,5-
di-t-butylbenzyl ester, etc.), aryl ester which may have
one or more suitable substituents (e.g. phenyl ester,
15 tolyl ester, t-butylphenyl ester, xylyl ester, mesityl
ester, cumenyl ester, salicyl ester, etc.), heterocyclic
ester (e.g. phthalidyl ester, etc.) and the like.

Suitable "lower alkylene" group may include straight
or branched one such as methylene, ethylene, trimethylene,
20 propylene, tetramethylene, hexamethylene, and the like,
in which the preferred one is C₁-C₂ alkylene and the
most preferred one is methylene.

Suitable "heterocyclic" group may include saturated
or unsaturated, monocyclic or polycyclic heterocyclic
25 group containing at least one hetero-atom such as an
oxygen, sulfur, nitrogen atom and the like.
And, especially preferable heterocyclic group may be
heterocyclic group such as

30 unsaturated 3 to 8-membered(more preferably
5 or 6-membered)heteromonocyclic group containing 1
to 4 nitrogen atom(s), for example, pyrrolol,
pyrrolinyl, imidazolyl, pyrazolyl, pyridyl and its
N-oxide, dihydropyridyl, pyrimidinyl, pyrazinyl,
35 pyridazinyl, triazolyl (e.g. 4H-1,2,4-triazolyl,

1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.),
tetrazolyl (e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.)
etc.;

5 saturated 3 to 8-membered(more preferably 5 or 6-
membered)heteromonocyclic group containing 1 to 4
nitrogen atom(s), for example, pyrrolidinyl,
imidazolidinyl, piperidino, piperazinyl, etc.;

10 unsaturated condensed heterocyclic group containing
1 to 4 nitrogen atom(s), for example, indolyl,
isoindolyl, indoliziny, benzimidazolyl, quinolyl,
isoquinolyl, indazolyl, benzotriazolyl, etc.;

15 unsaturated 3 to 8-membered(more preferably 5 or 6-
membered)heteromonocyclic group containing 1 to 2
oxygen atom(s) and 1 to 3 nitrogen atom(s), for
example, oxazolyl, isoxazolyl, oxadiazolyl (e.g.
1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl,
etc.), etc.;

20 saturated 3 to 8-membered (more preferably 5 or 6-
membered) heteromonocyclic group containing 1 to 2
oxygen atom(s) and 1 to 3 nitrogen atom(s), for
example, morpholinyl, sydnonyl, etc.;

25 unsaturated condensed heterocyclic group containing
1 to 2 oxygen atom(s) and 1 to 3 nitrogen atom(s),
for example, benzoxazolyl, benzoxadiazolyl, etc.;

30 unsaturated 3 to 8-membered (more preferably 5 or
6-membered) heteromonocyclic group containing 1 to
2 sulfur atom(s) and 1 to 3 nitrogen atom(s), for
example, thiazolyl, isothiazolyl, thiadiazolyl
(e.g. 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-
thiadiazolyl, 1,2,5-thiadiazolyl, etc.),
dihydrothiazinyl, etc.

35 saturated 3 to 8-membered (more preferably 5 or 6-
membered) heteromonocyclic group containing 1 to
2 sulfur atom(s) and 1 to 3 nitrogen atom(s), for
example, thiazolidinyl, etc.;

- unsaturated 3 to 8-membered (more preferably 5 or 6-membered) heteromonocyclic group containing 1 to 2 sulfur atom(s), for example, thienyl, dihydrodithiinyl, etc.;
- 5 unsaturated condensed heterocyclic group containing 1 to 2 sulfur atom(s) and 1 to 3 nitrogen atom(s), for example, benzothiazolyl, benzothiadiazolyl, etc.;
- unsaturated 3 to 8-membered (more preferably 5 or 6-membered) heteromonocyclic group containing an oxygen atom, for example, furyl, etc.;
- 10 unsaturated 3 to 8-membered (more preferably 5 or 6-membered) heteromonocyclic group containing an oxygen atom and 1 to 2 sulfur atom(s), for example, dihydro-oxathiinyl, etc.;
- 15 unsaturated condensed heterocyclic group containing an oxygen atom and 1 to 2 sulfur atom(s), for example, benzothienyl, benzodithiinyl, etc.;
- unsaturated condensed heterocyclic group containing an oxygen atom and 1 to 2 sulfur atom(s), for example, benzoxathiinyl, etc. and the like.

Thus defined heterocyclic group may optionally be substituted by one to ten, same or different, suitable substituent(s) such as:

- lower alkyl (e.g. methyl, ethyl, etc.);
- 25 lower alkoxy (e.g. methoxy, ethoxy, propoxy, etc.);
- lower alkylthio (e.g. methylthio, ethylthio, etc.);
- lower alkylamino (e.g. methylamino, etc.); cyclo-(lower)alkyl (e.g. cyclopentyl, cyclohexyl, etc.);
- 30 cyclo(lower)alkenyl (e.g. cyclohexenyl; cyclohexadienyl, etc.); hydroxy; halogen (e.g. chloro, bromo, etc.); amino; protected amino as aforementioned; cyano; nitro; carboxy; protected carboxy as aforementioned;
- sulfo; sulfamoyl; imino; oxo; amino(lower)alkyl (e.g. aminomethyl, aminoethyl, etc.); and the like.

- 35 Suitable "lower alkoxy carbonyl(lower)alkyl" group

42353D

42353D

may include ethoxycarbonylmethyl, propoxycarbonylmethyl, 1- or 2-ethoxycarbonylethyl, and the like.

Suitable "lower alkoxy carbonyl" moiety may include ethoxycarbonyl, propoxycarbonyl, and the like.

- 5 Suitable "halogen" may include chloro, bromo, iodo, and the like.

Suitable "conventional group which is capable to be replaced by the residue (-S-R^b) of the compound of the formula: HS-R^b" may include halogen as exemplified above.

10

(continued to the next page)

15

20

25


30

35


Suitable "trihalomethyl" may include trichloromethyl, and the like.

Particularly, the preferred embodiment of the symbols "R¹-A-", "R²" and "R³" of the object compounds (I) can be represented as follows.

The symbol " R^1-A- " can be represented by the formulae:-



R^1 is a group of the formula:



wherein R⁵ is amino or acylamino [more preferably lower alkanamido (e.g. formamido, acetamido, propionamido, etc.) or mono- or di- or trihalo(lower)alkanamido (e.g. chloroacetamido, dichloroacetamido, trifluoroacetamido, etc.)],

R⁶ is lower alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, pentyl, hexyl, etc.),

carboxy(lower)alkyl (e.g. carboxymethyl, 1-
or 2- carboxyethyl, 1- or 2- or 3- carboxypropyl,
etc.), or

esterified carboxy(lower)alkyl [more preferably

lower alkoxycarbonyl(lower)alkyl (e.g. methoxycarbonyl-

methyl, ethoxycarbonylmethyl, tert-butoxycarbonylmethyl,

tert-butoxycarbonyl ethyl, etc.) or mono- or di- or

tr-phenyl (lower) alkoxycarbonyl (lower) alkyl (e.g.

benzyl oxycarbonylmethyl, benzhydroyloxycarbonylmethyl,

benzhydrioxycarbonyl' etc.)) ; or


796

423530

4.8.0

② R¹-A- , in which

R^1 is a group of the formula:




wherein R⁵ is amino or acylamino (more preferably lower alkanamido (e.g. formamido, acetamido, propionamido, etc.)), and

A is methylene, aminomethylene, acylaminomethylene [more preferably lower alkoxy carbonylaminomethylene (e.g. methoxycarbonylaminomethylene, ethoxycarbonylaminomethylene, tert- butoxycarbonylaminomethylene, etc.)], hydroxymethylene or carbonyl; or



R_1 is a group of the formula:



wherein R^4 is lower alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, pentyl, etc.), and

A is aminomethylene or acylaminomethylene [more preferably lower alkoxy carbonylaminomethylene (e.g. methoxycarbonylaminomethylene, ethoxycarbonylaminomethylene, tert-butoxycarbonylaminomethylene, etc.)].

The symbol "R2" can be represented by:-

35 lower alkoxyethyl (e.g. methoxyethyl, ethoxyethyl,

962

423530

181

propoxymethyl, isopropoxymethyl, etc.); lower alkylthiomethyl (e.g. methylthiomethyl, ethylthiomethyl, propylthiomethyl, isopropylthiomethyl, etc.); or lower alkenylthiomethyl (e.g. vinylthiomethyl, allylthiomethyl, butenylthiomethyl, etc.).

The symbol "R³" can be represented by :-

carboxy or esterified carboxy [more preferably lower alkoxy-carbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, tert-butoxycarbonyl, etc.) or mono- or di- or triphenyl(lower)-alkoxycarbonyl (e.g. benzyloxycarbonyl, benzhydryloxycarbonyl, phenethylloxycarbonyl, etc.)].

The processes 1 to 13 for the preparation of the object compounds (I) of the present invention are explained in detail in the following.

(1) Process 1:

The compounds (I) or a salt thereof can be prepared by reacting the compound (II) or its reactive derivative at the amino group or a salt thereof with the compound (III) or its reactive derivative at the carboxy group or a salt thereof.

Suitable salts of the starting compounds (II) and (III) may include the same ones as illustrated for the compounds (I).


Suitable reactive derivative at the carboxy group of the compound (II) may include a conventional one, for example, a silyl derivative formed by the reaction of the compound (II) with a silyl compound such as bis-(trimethylsilyl)acetamide, trimethylsilylacetamide, etc.; isocyanate; isothiocyanate; Schiff's base or its tautomeric enamine type isomer formed by the reaction of the amino group with a carbonyl compound such as an aldehyde compound (e.g. acetaldehyde, isopentaldehyde, benzaldehyde, salicylaldehyde, phenylacetaldehyde, p-nitrobenzaldehyde, m-chlorobenzaldehyde, p-chlorobenzaldehyde, hydroxynaphthoaldehyde, furfural,

thiophenecarboaldehyde, etc.) or a ketone compound (e.g. acetone, methyl ethyl ketone, methyl isobutyl ketone, acetylacetone, ethyl acetoacetate, etc.), and the like.

5 Suitable reactive derivative of the compound (III) may include, for example, an acid halide, an acid anhydride, an activated amide, an activated ester, and the like, and preferably an acid chloride and acid bromide; a mixed acid anhydride with an acid such as substituted phosphoric acid (e.g. dialkylphosphoric acid, phenylphosphoric acid, diphenylphosphoric acid, dibenzylphosphoric acid, halogenated phosphoric acid, etc.), dialkylphosphorous acid, sulfurous acid, thiosulfuric acid, sulfuric acid, alkyl carbonate 15 (e.g. methyl carbonate, ethyl carbonate, propyl carbonate, etc.), aliphatic carboxylic acid (e.g. pivalic acid, pentanoic acid, isopentanoic acid, 2-ethylbutyric acid, trichloroacetic acid, etc.), aromatic carboxylic acid (e.g. benzoic acid, etc.); 20 a symmetrical acid anhydride; an activated acid amide with a heterocyclic compound containing imino function such as imidazole, 4-substituted imidazole, dimethylpyrazole, triazole or tetrazole; an activated ester (e.g. p-nitrophenyl ester, 2,4-dinitrophenyl 25 ester, trichlorophenyl ester, pentachlorophenyl ester, mesylphenyl ester, phenylazophenyl ester, phenyl thioester, p-nitrophenyl thioester, p-cresyl thioester, carboxymethyl thioester, pyridyl ester, piperidinyl ester, 8-quinolyl thioester, or an ester with a N-hydroxy 30 compound such as N,N-dimethylhydroxylamine, 1-hydroxy-2-(1H)-pyridone, N-hydroxysuccinimide, N-hydroxyphthalimide, 1-hydroxybenzotriazole, 1-hydroxy-6-chlorobenzotriazole, etc.), and the like.

Additionally, as a reactive derivative of the compound (III), wherein A is aminomethylene, the compound

of the following formula can also be used.



The suitable reactive derivative can optionally be selected from the above according to the kinds of the compounds (II) and (III) to be used practically.

This reaction can be carried out in the presence

of an organic or inorganic base such as alkali metal

(e.g. lithium, sodium, potassium, etc.), alkaline earth

metal (e.g. calcium, etc.), alkali metal hydride (e.g.

sodium hydride, etc.), alkaline earth metal hydride

(e.g. calcium hydride, etc.), alkali metal hydroxide

(e.g. sodium hydroxide, potassium hydroxide, etc.),

alkali metal carbonate (e.g. sodium carbonate, potassium

carbonate, etc.), alkali metal bicarbonate (e.g. sodium

bicarbonate, potassium bicarbonate, etc.), alkali metal

alkoxide (e.g. sodium methoxide, sodium ethoxide,

potassium tert-butoxide, etc.), trialkylamine (e.g.

triethylamine, etc.), pyridine compound (e.g. pyridine,

lutidine, picoline, etc.), quinoline, and the like.

In case that the compound (III) is used in a form

of the free acid or a salt in this reaction, the reaction

is preferably carried out in the presence of a condensing

agent such as a carbodiimide compound [e.g. N,N'-

dicyclohexylcarbodiimide, N-cyclohexyl-N'-morpholino-

ethylcarbodiimide, N-cyclohexyl-N'-(4-diethylamino-

cyclohexyl)carbodiimide, N,N'-diethylcarbodiimide,

N,N'-diisopropylcarbodiimide, N-ethyl-N'-(3-dimethyl-

aminopropyl)carbodiimide, etc.]; a ketenimine compound

(e.g. N,N'-carbonylbis(2-methylimidazole),

pentamethyleneketene-N-cyclohexylimine, diphenylketene-N-cyclohexylimine, etc.); an olefinic or acetylenic ether compounds (e.g. ethoxyacetylene, β -chlorovinyl-ethyl ether), a sulfonic acid ester of N-hydroxy-benzotriazole derivative [e.g. 1-(4-chlorobenzenesulfonyl-oxy)-6-chloro-1H-benzotriazole, etc.], a combination of trialkylphosphite or triphenylphosphine and carbon tetrachloride, disulfide or diazenedicarboxylate (e.g. diethyl diazenedicarboxylate, etc.), a phosphorus compound (e.g. ethyl polyphosphate, isopropyl polyphosphate, phosphoryl chloride, phosphorus trichloride, etc.), thionyl chloride, oxalyl chloride, N-ethylbenzoxazolium salt, N-ethyl-5-phenylisoxazolium-3-sulfonate, a reagent (referred to as so-called "Vijsmeier reagent") formed by the reaction of an amide compound such as dimethylformamide, N-methylformamide or the like with a halogen compound such as thionyl chloride, phosphoryl chloride, phosgene or the like.

The reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, acetone, dioxane, acetonitrile, chloroform, benzene, methylene chloride, ethylene chloride, tetrahydrofuran, ethyl acetate, N,N-dimethylformamide, pyridine, hexamethylphosphoramide, etc., or a mixture thereof.

Among these solvents, hydrophilic solvents may be used in a mixture with water.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

(2) Process 2:

The compound (I-b) or a salt thereof can be prepared by subjecting the compound (I-a) or a salt thereof to removal reaction of the amino-protective group in A¹.

Suitable method for this removal reaction may

include conventional one such as hydrolysis, reduction and the like.

(i) For Hydrolysis:

Hydrolysis is preferably carried out in the presence of an acid.

Suitable acid may be an inorganic acid (e.g.

hydrochloric acid, hydrobromic acid, sulfuric acid, etc.),

an organic acid (e.g. formic acid, acetic acid,

trifluoroacetic acid, propionic acid, benzenesulfonic

10 acid, p-toluenesulfonic acid, etc.), an acidic ion-

exchange resin and the like. In case that trifluoroacetic

acid is used in this reaction, the reaction is preferably

carried out in the presence of cation trapping agents

(e.g. anisole, etc.).

15 The acid suitable for this hydrolysis can be selected

according to the kinds of the protective group to be

removed, for example, this hydrolysis can preferably be

applied to the amino-protective group for A¹ such as

substituted or unsubstituted lower alkoxycarbonyl,

20 substituted or unsubstituted lower alkanoyl.

The hydrolysis is usually carried out in a con-

ventional solvent which does not adversely influence

the reaction such as water, methanol, ethanol, propanol,

tetrahydrofuran, N,N-dimethylformamide, dioxane or a

25 mixture thereof, and further the above-mentioned acids

can also be used as a solvent when they are in liquid.

The reaction temperature of this hydrolysis is not

critical, and the reaction is usually carried out under

cooling to at somewhat elevated temperature.

30

(ii) For Reduction:

Reduction is carried out in a conventional manner, including chemical reduction and catalytic reduction.

Suitable reducing agents to be used in chemical

35 reduction are a combination of a metal (e.g. tin, zinc,

iron, etc.) or metallic compound (e.g. chromium chloride, chromium acetate, etc.) and an organic or inorganic acid (e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, p-toluenesulfonic acid,

5 hydrochloric acid, hydrobromic acid, etc.).

Suitable catalysts to be used in catalytic reduc-

tion are conventional ones such as platinum catalysts

(e.g. platinum plate, spongy platinum, platinum black,

colloidal platinum, platinum oxide, platinum wire, etc.),

10 palladium catalysts (e.g. spongy palladium, palladium

black, palladium oxide, palladium on carbon, colloidal

palladium, palladium on barium sulfate, palladium on

barium carbonate, etc.), nickel catalysts (e.g. reduced

nickel, nickel oxide, Raney nickel, etc.), cobalt

15 catalysts (e.g. reduced cobalt, Raney cobalt, etc.),

iron catalysts (e.g. reduced iron, Raney iron, etc.),

copper catalysts (e.g. reduced copper, Raney copper,

Ullman copper, etc.) and the like.

The reduction manner can be selected according to

20 the kinds of the protective group to be removed, for

example, the chemical reduction can preferably be applied

to the amino-protective group for A¹ such as halo(lower)-

alkoxycarbonyl and the like, and catalytic reduction can

preferably be applied to that such as substituted or

25 unsubstituted ar(lower)alkoxycarbonyl, and the like.

The reduction is usually carried out in a conven-

tional solvent which does not adversely influence the

reaction such as water, methanol, ethanol, propanol,

N,N-dimethylformamide, or a mixture thereof.

30 Additionally, in case that the above-mentioned acids to

be used in chemical reduction are in liquid, they can

also be used as a solvent. Further, a suitable solvent

to be used in catalytic reduction may be the above-

mentioned solvent, and other conventional solvent such

35 as diethyl ether, dioxane, tetrahydrofuran, etc., or a

42353D

42353D

802

486

803

487

mixture thereof.

The reaction temperature of this reduction is not critical and the reaction is usually carried out under cooling to warming.

The present invention includes, within the scope of the invention, cases that the protected amino group in R^1 and/or the protected carboxy group for R^3 are transformed into free amino group and/or free carboxy group, respectively during the reaction.

10

(3) Process 3:

The compound (1-d) or a salt thereof can be prepared by subjecting the compound (1-c) or a salt thereof to removal reaction of the amino-protective group in R_a^1 .

This reaction is carried out by a conventional method such as hydrolysis, reduction, and the like.

The method of hydrolysis and reduction, and the reaction conditions (e.g. reaction temperature, solvent, etc.) are substantially the same as those illustrated for the removal reaction of the amino-protective group of the compound (1-a) in Process 2, and therefore are to be referred to said explanation.

The present invention includes, within the scope of the invention, cases that the protected amino group in A and/or the protected carboxy group(s) for R^3 and A are transformed into free amino group and/or free carboxy group, respectively during the reaction.

(4) Process 4:

The compound (1-f) or a salt thereof can be prepared by subjecting the compound (1-e) or a salt thereof to removal reaction of the carboxy-protective group for R_a^3 .

This reaction is carried out by a conventional method such as hydrolysis, reduction, and the like.

The method of hydrolysis and reduction, and the

804

423530

488

reaction conditions (e.g. reaction temperature, solvent, etc.) are substantially the same as those illustrated for the removal reaction of the amino-protective group of the compound (1-a) in Process 2, and therefore are to be referred to said explanation.

The present invention includes, within the scope of the invention, cases that the protected amino group(s) in R^1 and A and/or the protected carboxy group in A are transformed into free amino group(s) and/or a free carboxy group, respectively during the reaction.

10

(5) Process 5:

The compound (1-e) or a salt thereof can be prepared by introducing a carboxy-protective group into the compound (1-f) or a salt thereof.

The introducing agent of a carboxy-protective group to be used in this reaction may include a conventional esterifying agent such as an alcohol or its reactive equivalent (e.g. halide, sulfonate, sulfate, diazo compound, etc.), and the like.

This reaction can also be carried out in the presence of a base, and suitable examples thereof are the same as those given in the explanation of Process 1, and can preferably be carried out in the presence of metal iodide (e.g. sodium iodide, etc.).

This reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as N,N-dimethylformamide, tetrahydrofuran, dioxane, methanol, ethanol, etc., or a mixture thereof.

The reaction temperature is not critical, and the reaction is usually carried out under cooling to at somewhat elevated temperature.

In case that the alcohol is used as the introducing agent of a carboxy-protective group, the reaction can be carried out in the presence of a condensing agent as

805

423530

489

illustrated in Process 1.

(6) Process 6:

The compound (I-h) or a salt thereof can be prepared by reducing the compound (I-g) or a salt thereof.

The reduction can be carried out by a conventional method such as reduction using a reducing agent, catalytic reduction, and the like.

Suitable reducing agent may include a conventional one used for conversion of a carbonyl group to a hydroxymethyl group such as metal borohydride, for example, alkali borohydride (e.g. sodium borohydride, potassium borohydride, sodium cyanoborohydride, etc.), lithium aluminum hydride, etc.; diborane; and the like.

The catalyst to be used in the catalytic reduction may include the same ones as exemplified for the reduction in Process 2.

This reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, methanol, ethanol, tetrahydrofuran, dioxane; etc., or a mixture thereof.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

25 (7) Process 7:

The compound (I-i) or a salt thereof can be prepared by reacting the compound (IV) or a salt thereof with the compound (V).

Suitable salts of the starting compound (IV) may include the same salts with a base for the compounds (I).

This reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as ethyl acetate, methylene chloride, chloroform, carbon tetrachloride, tetrahydrofuran, dioxane, water, etc., or a mixture thereof.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

(8) Process 8:

The compound (I-l) or a salt thereof can be prepared by subjecting the compound (I-k) or a salt thereof to removal reaction of the carboxy-protective group in A⁶.

This reaction is carried out by a conventional method such as hydrolysis, reduction, and the like.

The method of hydrolysis and reduction, and the reaction conditions (e.g. reaction temperature, solvent, etc.) are substantially the same as those illustrated for the removal reaction of the amino-protective group of the compound (I-a) in Process 2, and therefore are to be referred to said explanation.

The present invention includes, within the scope thereof, cases that the protected amino group in R¹ and/or the protected carboxy group in R³ are transformed into free amino group and/or free carboxy group, respectively during the reaction.

(continued to the next page)

(9.) Process 9:

The compound (I-n) or a salt thereof can be prepared by subjecting the compound (I-m) or a salt thereof to removal reaction of the amino- and carboxy-protective groups in R₃.

This reaction is carried out by a conventional method such as hydrolysis, reduction, and the like.

The method of hydrolysis and reduction, and the reaction conditions (e.g. reaction temperature, solvent, etc.) are substantially the same as those illustrated for removal reaction of the amino-protective group of the compound (I-a) in Process 2, and therefore are to be referred to said explanation.

In this reaction, the amino- and carboxy-protective groups can be removed separately or at a time.

(10) Process 10:

The compound (I-p) or a salt thereof can be prepared by subjecting the compound (I-o) or a salt thereof to removal reaction of the amino- and carboxy-protective groups in A⁸.

This reaction is carried out by a conventional method such as hydrolysis, reduction, and the like.

The method of hydrolysis and reduction, and the reaction conditions (e.g. reaction temperature, solvent, etc.) are substantially the same as those illustrated for removal reaction of the amino-protective group of the compound (I-a) in Process 2, and therefore are to be referred to said explanation.

In this reaction, the amino- and carboxy-protective groups can be removed separately or at a time.

(11) Process 11:

The compound (I-k) or a salt thereof can be prepared by introducing a carboxy-protective group into

the compound (I-l) or a salt thereof.

This reaction is carried out by substantially the same method as that illustrated for introducing the carboxy-protective group into the compound (I-f) in Process 5, and therefore, the reaction conditions (e.g. reaction temperature, solvent, etc.) are to be referred to said explanation.

(12) Process 12:

The compound (I-r) or a salt thereof can be prepared by reacting the compound (I-q) or a salt thereof with the compound (VII).

This reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as tetrahydrofuran, dioxane, water, etc., or a mixture thereof.

The reaction temperature is not critical, and the reaction is usually carried out at ambient temperature to under heating.

(13) Process 13:

The compound (I-s) or a salt thereof can be prepared by reacting the compound (I-r) or a salt thereof with a base.

Suitable base used in this Process may include the same ones as those exemplified in Process 1.

This reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, methanol, ethanol, etc., or a mixture thereof.

The reaction temperature is not critical, and the reaction is usually carried out under cooling to warming.

The object compounds (I) obtained according to the Processes 1 to 13 as explained above can be isolated and purified in a conventional manner, for example, extraction,

precipitation, fractional crystallization, recrystallization, chromatography, and the like.

Processes A to F for the preparation of the starting compounds (II-d), (II-g), (III-b) to (III-e), (III-g), (III-h), (IV) and (XIV-c) are explained in detail in the following.

Process A-1:

The compound (II-b) or a salt thereof can be prepared by reacting the compound (II-a) or a salt thereof with the compound (VIII) or its reactive derivative at the mercapto group.

Suitable salts of the compounds (II-a) and (II-b) may include the same salts with a base as exemplified for the compounds (I).

Suitable "reactive derivative at the mercapto group" of the compound (VIII) may include salts with a base as exemplified for the compounds (I).

This reaction is preferably carried out in the presence of a base and suitable examples thereof are the same as those given in the explanation of Process 1.

The reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as N,N-dimethylformamide, dimethylsulfoxide, methanol, ethanol, chloroform, etc., or a mixture thereof.

The reaction temperature is not critical and the reaction is usually carried out at ambient temperature to under warming.

30 Process A-2:

The compound (II-c) or a salt thereof can be prepared by reducing the compound (II-b) or a salt thereof.

Suitable salts of the compound (II-c) may include the same salts with a base as exemplified for the compounds (I).

The reducing agent to be used in this reaction may include a conventional one used for conversion of a sulfinyl group to a thio group such as phosphorus halide (e.g. phosphorus trichloride, phosphorus pentachloride, etc.); stannous halide (e.g. stannous chloride, etc.); silicon halide (e.g. silicon tetrachloride, etc.); excess amount of acid halide such as lower alkanoyl halide (e.g. acetyl bromide, acetyl chloride, etc.); combination of alkali metal halide (e.g. sodium iodide, etc.) and acid anhydride such as halo(lower)alkanoic anhydride (e.g. trifluoroacetic anhydride, etc.); and the like.

The reaction is usually carried out in the presence of an acid scavenger such as lower alkene (e.g. 2-methyl 2-butene, etc.), lower alkylene oxide (e.g. ethylene oxide, propylene oxide, etc.) and the like.

The reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as chloroform, methylene chloride, tetrahydrofuran, benzene, etc., or a mixture thereof.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

Process A-3:

The compound (II-d) or a salt thereof can be prepared by subjecting the compound (II-c) or a salt thereof to removal reaction of the amino-protective group for R^a.

Suitable salts of the compound (II-d) may include the same ones as exemplified for the compounds (I).

Suitable method for this removal reaction may include conventional one such as a combined method comprising iminohalogenation and iminoetherification, optionally followed by hydrolysis, and the like.

The first and second steps of this method are preferably carried out in an anhydrous solvent. Suitable solvent for the first step (i.e. iminohalogenation) is

an aprotic solvent such as methylene chloride, chloroform, diethyl ether, tetrahydrofuran, dioxane, etc., and for the second step (i.e. iminoetherification) is usually the same as those in the above first step. These two steps are usually conducted under cooling. These two steps and the last step (i.e. hydrolysis step) are most preferably conducted in one-batch system.

Suitable iminohalogenating agents include a halogenating agent such as phosphorus halo compound (e.g. phosphorus trichloride, phosphorus pentachloride, phosphorus tribromide, phosphorus pentabromide, phosphorus oxychloride, etc.), thionyl chloride, phosgene, and the like.

Suitable iminoetherifying agent may be an alcohol such as an alcohol (e.g. methanol, ethanol, propanol, isopropanol, butanol, etc.) or the corresponding alcohol having alkoxy (e.g. 2-methoxyethanol, 2-ethoxyethanol, etc.), and alkoxide of metal such as alkali metal, alkaline earth metal (e.g. sodium methoxide, potassium ethoxide, magnesium ethoxide, lithium methoxide, etc.), and the like.

Thus obtained reaction product is, if necessary, hydrolyzed in a conventional manner.

The hydrolysis is preferably carried out at ambient temperature or under cooling, and proceeds simply pouring the reaction mixture into water or a hydrophilic solvent such as alcohol (e.g. methanol, ethanol, etc.) moistened or admixed with water, and if necessary, with addition of an acid or base as exemplified in Processes 1 and 2.

Process B-1:

The compound (III-b) can be prepared by reacting the compound (III-a) with the compound (IX) or a salt thereof.

Suitable salt of the compound (IX) may include the same one as exemplified for the compounds (I).

This reaction is preferably carried out in the presence of a base as exemplified in Process 1.

This reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, dioxane, tetrahydrofuran, etc., or a mixture thereof.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

Process B-2:

The compound (III-c) or a salt thereof can be prepared by reducing the compound (III-b).

Suitable salts of the compound (III-c) may include the same acid addition salt as exemplified for the compounds (I).

The reduction can be carried out by a conventional method such as chemical reduction, catalytic reduction, and the like.

The method of chemical reduction and catalytic reduction, and the reaction conditions (e.g. reaction temperature, solvent, etc.) are substantially the same as those illustrated for Process 2, and therefore are to be referred to said explanation.

Process B-3:

The compound (III-d) can be prepared by introducing an amino-protective group into the compound (III-c) or a salt thereof.

The introducing agent of an amino-protective group to be used in this reaction may include a conventional acylating agent such as the corresponding acid to the acyl group as aforementioned or its reactive derivative

423530

423530

(e.g. acid halide, acid anhydride, etc.), 2-lower alkoxy-carbonyloxyimino-2-phenylacetone nitrile (e.g. 2-tert-butoxy-carbonyloxyimino-2-phenylacetone nitrile, etc.), alkyl ketone substituted by lower alkoxy-carbonyl (e.g. lower alkyl acetoacetate, for example, methyl acetoacetate, etc., etc.), and the like.

This reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, methanol, ethanol, propanol, tetrahydrofuran, dioxane, etc., or a mixture thereof.

This reaction is preferably carried out in the presence of a base, and suitable examples thereof are the same as those given in the explanation of Process 1.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

Process B-4:

The compound (III-e) or a salt thereof can be prepared by subjecting the compound (III-d) to removal reaction of the carboxy-protective group.

Suitable salts of the compound (III-e) may include the same salts with a base as exemplified for the compounds (I).

This reaction is carried out by a conventional method such as hydrolysis, reduction, and the like.

The method of hydrolysis and reduction, and the reaction conditions (e.g. reaction temperature, solvent, etc.) are substantially the same as those illustrated for the removal reaction of the amino-protective group in Process 2, and therefore are to be referred to said explanation. Additionally, hydrolysis can be carried out in the presence of a base, and suitable examples thereof are the same as those in the explanation of Process 1.

Process B-5:

The compound (III-g) or a salt thereof can be prepared by subjecting the compound (III-f) or a salt thereof to removal reaction of the carboxy-protective group.

Suitable salts of the compound (III-f) may include the same acid addition salts as exemplified above, and those of the compound (III-g) may include the same salts as exemplified for the compounds (I).

The reaction is substantially the same as Process B-4, and therefore, the reaction method, reaction conditions (e.g. reaction temperature, solvent, etc.) are to be referred to said explanation.

Process B-6

The compound (III-e) or a salt thereof can be prepared by introducing an amino-protective group into the compound (III-g) or a salt thereof.

This reaction is substantially the same as Process B-3, and therefore, the reaction method and reaction conditions (e.g. reaction temperature, solvent, etc.) are to be referred to said explanation.

Process C:

The compound (IV) or a salt thereof can be prepared by reacting the compound (IV-a) or its reactive derivative at the amino group or a salt thereof with the compound (X) or its reactive derivative at the carboxy group or a salt thereof.

Suitable salts of the compound (IV-a) may include the same ones as exemplified for the compounds (I), and those of the compound (X) may include the same salts with a base as exemplified above.

The reaction is substantially the same method as Process 1, and accordingly, the method, reaction conditions (e.g. reaction temperature, solvent, base, etc.)

are to be referred to said explanation.

In this process, carbonyl equivalents, for example, acetal of the compound (X), wherein A₅ is lower alkylene having oxo, can also be used in this reaction and such acetal can easily be transformed into the oxo group by a conventional method (e.g. hydrolysis, etc.) after the reaction.

10

(continued to the next page)

15

Process D-1:

The compound (II-f) or a salt thereof can be prepared by reacting the compound (II-e) or a reactive derivative at the carboxy group or a salt thereof with lower alkanol substituted by protected amino and protected carboxy groups (XII).

Suitable salts of the compounds (II-e) and (II-f) may include the same ones as exemplified for the compounds (I).

10

Suitable reactive derivative at the carboxy group of the compound (II-e) may include the same ones as the compound (III) in Process 1.

This reaction is carried out by substantially the same method as that illustrated for introducing the carboxy-protective group into the compound (I-f) in Process 5, and therefore, the reaction conditions (e.g. reaction temperature, solvent, etc.) are to be referred to said explanation.

15

Process D-2:

The compound (II-g) or a salt thereof can be prepared by subjecting the compound (II-f) or a salt thereof to removal reaction of the amino-protective group for R^a.

Suitable salt of the compound (II-g) may include the same ones as exemplified for the compounds (I).

25

This reaction is carried out by substantially the same method as that illustrated for removal reaction of the amino-protective group of the compound (II-c) in Process A-3, and therefore, the reaction conditions (e.g. reaction temperature, solvent, etc.) are to be referred to said explanation.

30

Process E-1:

The compound (III-1) can be prepared by reacting the compound (III-h) with hydroxylamine or a salt thereof.

35

42353D

816

817

42353D

501

500

35

25

20

Suitable salt of hydroxylamine may include the same acid addition salt as exemplified for the compounds (I).

In case that the salt of hydroxylamine is used as the reagent, the reaction can usually be carried out in the presence of a base such as those illustrated in Process 1.

The reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as methanol, ethanol, etc., or a mixture thereof.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

Process E-2:

The compound (III-j) can be prepared by reacting the compound (III-i) with the compound (XIII) or a reactive derivative at the carboxy group or a salt thereof.

Suitable salt of the compound (XIII) may include the same salt with a base as exemplified for the compounds (I).

Suitable reactive derivative at the carboxy group of the compound (XIII) may include the same ones as illustrated for the compound (II) in Process 1.

This reaction is carried out by substantially the same method as Process 1, and therefore, the reaction conditions (e.g. reaction temperature, solvent, etc.) are to be referred to said explanation.

Process E-3:

The compound (III-k) or a salt thereof can be prepared by reacting the compound (III-j) or a salt thereof with ammonia.

Suitable salt of the compound (III-k) may include the same acid addition salt as exemplified for the compounds (I).

This reaction can be carried out in the absence of

or in the presence of a solvent which does not adversely influence the reaction such as dioxane, etc., and the reaction is usually carried out in the absence of a solvent.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

In case that the compound (III-k) is one of the geometrical isomers, it can be transformed into the other isomer in a conventional manner.

Process E-4:

The compound (III-l) or a salt thereof can be prepared by subjecting the compound (III-k) or a salt thereof to removal reaction of the carboxy-protective group for R.

Suitable salt of the compound (III-l) may include the same ones as exemplified for the compounds (I).

This reaction is carried out by a conventional method such as hydrolysis, reduction, and the like.

The method of hydrolysis and reduction, and the reaction conditions (e.g. reaction temperature, solvent, etc.) are substantially the same as those illustrated for the removal reaction of the amino-protective group of the compound (I-a) in Process 2, and therefore are to be referred to said explanation.

Process F-1:

The compound (XIV-b) can be prepared by reacting the compound (XIV-a) or a reactive derivative at the hydroxy group with N-hydroxyphthalimide.

Suitable reactive derivative at the hydroxy group may include halide such as chloride, bromide, and the like.

This reaction is preferably carried out in the presence of a base as exemplified in Process 1.

In case that the compound (XIV-a) is used in a free

form, the reaction can usually be carried out in the presence of a condensing agent as exemplified in Process 1.

This reaction is usually carried out in a conventional solvent which does not adversely influence the reaction such as tetrahydrofuran, N,N-dimethylformamide, etc., or a mixture thereof.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

Process F-2:

The compound (XIV-c) or a salt thereof can be prepared by subjecting the compound (XIV-b) to removal reaction of the phthaloyl group.

Suitable salt of the compound (XIV-c) may include the same acid addition salt as exemplified for the compounds (I).

This reaction is carried out by a conventional method such as hydrolysis, and the like.

The method of hydrolysis, and the reaction conditions (e.g. reaction temperature, solvent, etc.) are substantially the same as those illustrated for removal reaction of the amino-protective group of the compound (I-a) in Process 2, and therefore are to be referred to said explanation.

The starting compounds (II-d), (II-g), (III-b) to (III-e), (III-f), (IV) and (XIV-c) thus prepared can be isolated in a conventional manner as mentioned for the object compounds of the present invention.

It is to be noted that, in the aforementioned reactions in Processes 1 to 13 and A to F or the post-treatment of the reaction mixture therein, in case that the starting or object compounds possess an optical and/or geometrical isomer(s), it may occasionally be transformed into the other optical and/or geometrical

isomer(s), and such cases are also included within the scope of the present invention.

In case that the object compounds (I) have a free carboxy group or free amino group at the 4th or 7th position thereof, it may be transformed into its pharmaceutically acceptable salts by a conventional method.

The object compounds (I) and the pharmaceutically acceptable salts thereof of the present invention are novel and exhibit high antimicrobial activity, inhibiting the growth of a wide variety of pathogenic microorganisms including Gram-positive and Gram-negative microorganisms and are useful as antimicrobial agents, especially for oral administration.

Now in order to show the utility of the object compounds (I), the test data on the in vitro antimicrobial activity of some representative compounds (I) of this invention are shown in the following.

Test: In vitro Antimicrobial Activity.

Test Compounds

- | | |
|-------|---|
| No. 1 | 7-[2-(2-Aminothiazol-4-yl)-DL-glycolamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (hereinafter referred to as Compound A) |
| No. 2 | 7-[2-(3-Methanesulfonamidophenyl)-D-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid trifluoroacetate (hereinafter referred to as Compound B) |
| No. 3 | 7-[2-(3-Methanesulfonamidophenyl)-D-glycinamido]-3-allylthiomethyl-3-cephem-4-carboxylic acid trifluoroacetate (hereinafter referred to as Compound C) |
| No. 4 | 7-[2-(3-Methanesulfonamidophenyl)-D-glycinamido]-3-methoxymethyl-3-cephem-4-carboxylic acid (hereinafter referred to as Compound D) |

Test Method

In vitro Antimicrobial activity was determined by the two-fold agar-plate dilution method as described below.

One loopful of an overnight culture of each test strain in Trypticase-soy broth (approximately 10^8 viable cells per ml) was streaked on heart infusion agar (HI-agar) containing graded concentrations of antimicrobial agents, and the minimal inhibitory concentration (MIC) was expressed in term of $\mu\text{g/ml}$ after incubation at 37°C for 20 hours.

Test Results 1MIC ($\mu\text{g/ml}$)

Microorganisms Test compounds	Staphylococcus aureus 209P JC-1	Batillus subtilis ATCC 6633
A	1.56	0.78
B	1.56	0.10
C	3.13	0.78
D	1.56	0.39

(continued to the next page)

35

42353D

822

506

823

507

For therapeutic administration, the object compounds (I) and the pharmaceutically acceptable salts thereof of the present invention are used in the form of conventional pharmaceutical preparation which contains said compound, as active ingredients, in admixture with pharmaceutically acceptable carriers such as an organic or inorganic solid or liquid excipient which is suitable for oral, parenteral and external administration. The pharmaceutical preparations may be in solid form such as tablet, granule, powder, capsule, or liquid form such as solution, suspension, syrup, emulsion, lemonade and the like.

If needed, there may be included in the above preparations auxiliary substances, stabilizing agents, wetting agents and other commonly used additives such as lactose, magnesium stearate, terra alba, sucrose, corn starch, talc, stearic acid, gelatin, agar, pectin, peanut oil, olive oil, cacao butter, ethyleneglycol and the like.

While the dosage of the compounds (I) may vary from and also depend upon the age, conditions of the patient, a kind of diseases, a kind of the compounds (I) to be applied, etc. In general, amounts between 1 mg and about 4,000 mg or even more per day may be administered to a patient. An average single dose of about 50 mg, 100 mg, 250 mg, 500 mg, 1000 mg, 2000 mg of the object compounds (I) of the present invention may be used in treating diseases infected by pathogenic microorganisms.

The following examples are given for the purpose of illustrating the present invention.

(continued to the next page)

35

42353D

Preparation of the starting compoundsPreparation 1

To a solution of benzhydryl 7-(2-phenylacetamido)-3-chloromethyl-3-cephem-4-carboxylate-1-oxide (25 g) in N,N-dimethylformamide (150 ml) were added triethylamine (6.42 ml) and 2-propene-1-thiol (8.0 ml), and the mixture was stirred at 25°C for 3 hours.

The reaction mixture was poured into a saturated aqueous solution of sodium chloride (1.5 l), followed by collecting the precipitated solid by filtration, which was washed with water and diisopropyl ether, and then dried to give benzhydryl 7-(2-phenylacetamido)-3-allylthiomethyl-3-cephem-4-carboxylate-1-oxide (24.2 g).

15 I.R. (Nujol): 1775, 1715, 1644, 1170, 1030 cm^{-1}

NMR δ ppm (DMSO- d_6): 3.00 (2H, d, J=7Hz), 3.6 (6H, m), 5.0 (1H, d, J=5Hz), 5.90 (1H, dd, J=5Hz, 8Hz), 4.8-5.6 (3H, m), 7.00 (1H, s), 7.4 (15H, s), 8.40 (1H, d, J=8Hz)

Preparation 2

Benzhydryl 7-(2-phenylacetamido)-3-methylthiomethyl-3-cephem-4-carboxylate-1-oxide (13.7 g) was obtained by reacting benzhydryl 7-(2-phenylacetamido)-3-chloromethyl-3-cephem-4-carboxylate-1-oxide (15 g) with 30% methanolic methanethiol (15 ml) in substantially the same manner as that of Preparation 1.

I.R. (Nujol): 3300, 1775, 1710, 1650, 1172, 1027 cm^{-1}

NMR δ ppm (DMSO- d_6): 1.80 (3H, s), 3.3-4.0 (6H, m), 5.02 (1H, d, J=5Hz), 5.93 (1H, dd, J=5Hz, 8Hz), 7.02 (1H, s), 7.50 (15H, s), 8.40 (1H, d, J=8Hz)

Preparation 3

Benzhydryl 7-(2-phenylacetamido)-3-ethylthiomethyl-3-cephem-4-carboxylate-1-oxide (14.1 g) was obtained

42353D

824

508

by reacting benzhydryl 7-(2-phenylacetamido)-3-chloromethyl-3-cephem-4-carboxylate-1-oxide (15 g) with ethanethiol (4.05 ml) in substantially the same manner as that of Preparation 1.

5 I.R. (Nujol): 3280, 1776, 1708, 1647, 1172, 1015 cm^{-1}

NMR δ ppm (DMSO- d_6): 0.95 (3H, t, J=7Hz), 2.28 (2H, q, J=7Hz), 3.5-4.0 (6H, m), 5.02 (1H, d, J=5Hz), 5.93 (1H, dd, J=5Hz, 8Hz), 7.02 (1H, s), 7.5 (15H, s), 8.43 (1H, d, J=8Hz)

Preparation 4

To a solution of benzhydryl 7-(2-phenylacetamido)-3-allylthiomethyl-3-cephem-4-carboxylate-1-oxide (26 g) in methylene chloride (500 ml) was added dropwise phosphorus trichloride (20 ml) at 5°C with stirring, and the stirring was continued for an hour. The mixture was poured into a mixture of methylene chloride (200 ml) and water (400 ml), followed by separating out the organic layer, which was washed twice with an aqueous solution of sodium chloride (200 ml) and dried over anhydrous magnesium sulfate. After removal of the solvent, the residue was pulverized with diisopropyl ether to give benzhydryl 7-(2-phenylacetamido)-3-allylthiomethyl-3-cephem-4-carboxylate (22 g).

I.R. (Nujol): 1770, 1715, 1650 cm^{-1}

NMR (DMSO- d_6): 3.0 (2H, d, J=7Hz), 3.6 (6H, m), 5.00 (1H, d, J=5Hz), 5.67 (1H, dd, J=5Hz, 8Hz), 4.8-5.5 (3H, m), 6.90 (1H, s), 7.40 (15H, m), 9.10 (1H, d, J=8Hz)

Preparation 5

To a solution of benzhydryl 7-(2-phenylacetamido)-3-methylthiomethyl-3-cephem-4-carboxylate-1-oxide

42353D

825

509

(15.1 g) in methylene chloride (150 ml) was added 2-methyl-2-butene (5.7 ml), followed by adding dropwise acetyl bromide (5.2 ml) under ice-cooling and stirring for half an hour. After addition of water, the mixture was adjusted to pH about 5 with an aqueous solution of sodium bicarbonate, washed three times with an aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate and then evaporated under reduced pressure to give benzhydryl 7-(2-phenylacetamido)-3-methylthiomethyl-3-cephem-4-carboxylate (13.5 g).

I.R. (Nujol): 3380, 1785, 1715, 1652 cm^{-1}

NMR δ ppm ($\text{DMSO}-d_6$): 1.83 (3H, s), 3.60 (4H, broad s), 3.66 (2H, broad s), 5.23 (1H, d, J=5Hz), 5.75 (1H, dd, J=5Hz, 8Hz), 6.97 (1H, s), 7.43 (15H, s), 9.17 (1H, d, J=8Hz)

Preparation 6

Benzhydryl 7-(2-phenylacetamido)-3-ethylthiomethyl-3-cephem-4-carboxylate (23 g) was obtained by reacting benzhydryl 7-(2-phenylacetamido)-3-ethylthiomethyl-3-cephem-4-carboxylate-1-oxide (30 g) with acetyl bromide (10.2 ml) in the presence of 2-methyl-2-butene (11.1 ml) in substantially the same manner as that of Preparation 5.

I.R. (Nujol): 3300, 1772, 1701, 1650 cm^{-1}

NMR δ ppm ($\text{DMSO}-d_6$): 1.00 (3H, t, J=7Hz), 2.33 (2H, q, J=7Hz), 3.56 (6H, broad s), 5.17 (1H, d, J=5Hz), 5.76 (1H, dd, J=5Hz, 8Hz), 7.00 (1H, s), 9.13 (1H, d, J=8Hz)

Preparation 7

To a suspension of phosphorus pentachloride (16.1 g) and pyridine (6.3 ml) in methylene chloride (100 ml) were added benzhydryl 7-(2-phenylacetamido)-

3-allylthiomethyl-3-cephem-4-carboxylate (22 g) and methylene chloride (100 ml) at 5°C, and the mixture was stirred at the same temperature for an hour.

After cooling to -20°C, methanol (10 ml) was added thereto, followed by stirring at -10°C for half an hour. To this mixture was added water (10 ml) and stirred for 10 minutes. To the separated methylene chloride was added an aqueous solution of sodium bicarbonate until the pH value of the aqueous solution became 5.0, and the mixture was shaken. After separating out the organic layer, it was washed with an aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate and evaporated to dryness. The residue obtained was pulverized with diisopropyl ether to give benzhydryl 7-amino-3-allylthiomethyl-3-cephem-4-carboxylate (6.5 g).

I.R. (Nujol): 1770, 1710 cm^{-1}

NMR δ ppm ($\text{DMSO}-d_6$): 2.93 (2H, d, J=7Hz), 3.3-3.7 (4H, m), 5.00 (1H, d, J=5Hz), 5.60 (1H, d, J=5Hz), 4.5-5.6 (3H, m), 6.91 (1H, s), 7.5 (10H, m)

Preparation 8

Benzhydryl 7-amino-3-methylthiomethyl-3-cephem-4-carboxylate (5.0 g) was obtained by reacting benzhydryl 7-(2-phenylacetamido)-3-methylthiomethyl-3-cephem-4-carboxylate (13.5 g) with phosphorus pentachloride (7.74 g), pyridine (3 ml) and methanol (100 ml) in substantially the same manner as that of Preparation 7.

I.R. (Nujol): 1765, 1725 cm^{-1}

NMR δ ppm ($\text{DMSO}-d_6$): 1.81 (3H, s), 3.52 (2H, broad s), 3.60 (2H, broad s), 4.83, 5.13 (2H, ABq, J=5Hz), 6.97 (1H, s), 7.40 (10H, s)

Preparation 9

Benzhydryl 7-amino-3-ethylthiomethyl-3-cephem-4-

carboxylate (10.0 g) was obtained by reacting benzhydryl 7-(2-phenylacetamido)-3-ethylthiomethyl-3-cephem-4-carboxylate (23.0 g) with phosphorus pentachloride (12.90 g), pyridine (5.0 ml) and methanol (165 ml) in substantially the same manner as that of Preparation 7.

I.R. (Nujol): 1770, 1720 cm^{-1}

¹H-NMR (DMSO- d_6): 0.96 (3H, t, J=7Hz), 2.30 (2H, q, J=7Hz), 3.50 (2H, broad s), 3.60 (2H, broad s), 4.80, 5.17 (2H, ABq, J=5Hz), 7.00 (1H, s), 7.43 (10H, s)

(continued to the next page)

Preparation 10

To a solution of diketene (1.3 ml) in methylene chloride (10 ml) was added dropwise a solution of bromine (1.24 g) in methylene chloride (10 ml) at -30°C with stirring, and the stirring was continued at -20°C for half an hour to prepare a solution of

4-bromoacetoacetyl bromide. This solution was added dropwise to a solution of benzhydryl 7-amino-3-methylthiomethyl-3-cephem-4-carboxylate (4.44 g) and trimethylsilylacetylamide (5.46 g) in methylene chloride (100 ml) at -30 to -20°C over a period of 5 minutes with stirring, and the stirring was continued at -10°C for half an hour. After addition of water, the resultant mixture was adjusted to pH 7.5 with an aqueous solution of sodium bicarbonate, followed by separating out the organic layer, which was washed with water and an aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and then evaporated to dryness to give benzhydryl 7-(4-bromoacetoacetamido)-3-methylthiomethyl-3-cephem-4-carboxylate (6.0 g).

IR (Nujol) : 1770, 1710, 1625 cm^{-1}

NMR δ ppm (DMSO- d_6) : 1.77 (3H, s), 3.6 (6H, m), 4.33 (2H, s), 5.15 (1H, d, J=4Hz), 5.73 (1H, dd, J=4Hz, 8Hz), 6.86 (1H, s), 7.3 (10H, m), 9.1 (1H, d, J=8Hz)

Preparation 11

To a solution of ethyl 2-(2-formamidothiazol-4-yl)-2-methoxyiminoacetic acid (syn isomer) (19 g) in methanol (200 ml) were added 50% formic acid (200 ml) and zinc (29 g), and the mixture was stirred at 5 to 10°C for 6 hours. After filtration, the reaction mixture was evaporated, followed by dissolving the residue in water (150 ml). The resultant aqueous solution was adjusted to pH 6.5 with 4N aqueous solution of sodium hydroxide, followed by addition of ethanol (150 ml),

2-tert-butoxycarbonyloxyimino-2-phenylacetoneitrile (18.2 g) and triethylamine (8.0 g): After stirring at ambient temperature for 24 hours, the reaction mixture was filtered, followed by removal of the organic solvent. The remained aqueous solution was washed with ethyl acetate, adjusted to pH 4 with 10% hydrochloric acid and then extracted with ethyl acetate. The extract was washed with an aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate and evaporated to dryness under reduced pressure to give a residue, which was washed with diethyl ether to obtain N-tert-butoxycarbonyl-2-(2-formamidothiazol-4-yl)glycine (3.3 g).

IR (Nujol) : 3250, 3180, 1720, 1700, 1670, 1640, 1540, 1510 cm^{-1}

NMR δ ppm ($\text{DMSO}-d_6$) : 1.40 (9H, s), 5.18 (1H, d, J=8Hz), 7.17 (1H, s), 8.43 (1H, s)

(continued to the next page)

830

42353D

514

35

Preparation 12

Bromoacetic acid (10.45 g) was dissolved in methanol (30 ml). To the solution was added an equivalent volume of diphenyl diazomethane in ethyl acetate at 45°C, and the reaction mixture was stirred at the same temperature for an hour. The solution was washed with 5% aqueous sodium bicarbonate and a saturated aqueous sodium chloride, and then dried over magnesium sulfate. The solution was evaporated in vacuo to give an oily product. This oil was dissolved in N,N-dimethylformamide (60 ml). To the solution was added N-hydroxyphthalimide (11.7 g) and triethylamine (15.1 ml), and the reaction mixture was stirred at ambient temperature for an hour. The resultant mixture was poured into a saturated aqueous sodium chloride (500 ml). The precipitates were collected by filtration, washed with water, and then dissolved in methylene chloride (700 ml). The solution was washed with a saturated aqueous sodium chloride, dried over magnesium sulfate, and then evaporated under reduced pressure to give benzhydryl 2-phthalimidoxyacetate (20.4 g), mp 173-175°C.

IR (Nujol) : 1754, 1730 cm^{-1}

NMR δ ppm (CDCl_3 , δ) : 4.93 (2H, s), 7.0 (1H, s), 7.3 (10H, s), 7.73 (4H, s).

Preparation 13

To a solution of benzhydryl 2-phthalimidoxyacetate (10 g) in methylene chloride (100 ml) was added a solution of hydrazine hydrate (6.08 g) in methanol (7 ml). The reaction mixture was stirred at ambient temperature for an hour. The precipitates were collected by filtration and washed with methylene chloride. The filtrate and the washings were combined, adjusted to pH 7.0 with conc. hydrochloric acid, and washed with a saturated aqueous sodium chloride, and then dried over magnesium sulfate.

831

42353D

515

35

The solution was evaporated in vacuo to give benzhydryl 2-aminoxyacetate (6.0 g).

IR (film): 3320, 1750 cm^{-1}

NMR δ ppm (CDCl_3 , δ): 4.33 (2H, s), 5.86

(2H, broad s), 7.00 (1H, s), 7.3 (10H, s)

Preparation 14

To a suspension of (2-formamidothiazol-4-yl)-glyoxylic acid (6.0 g) in water (60 ml) and pyridine (6 ml) was added a solution of benzhydryl 2-aminoxyacetate (9.0 g) in tetrahydrofuran (40 ml).

The reaction mixture was stirred at ambient temperature for 3 hours. To the resultant solution was added ethyl acetate (200 ml). The separated organic

layer was washed with 5% hydrochloric acid (100 ml), a saturated aqueous sodium bicarbonate and a saturated aqueous sodium chloride, and then dried over magnesium sulfate. The solvent was distilled off to give 2-(2-formamidothiazol-4-yl)-2-benzhydryloxyacarbonylmethoxyaminoacetic acid (syn isomer) (13.0 g), mp 143-151°C.

IR (Nujol): 3150, 1733, 1692 cm^{-1}

NMR δ ppm (DMSO-d_6): 5.0 (2H, broad s), 6.97 (1H, s), 7.40 (20H, m), 7.56 (1H, s), 8.60 (1H, s), 12.77 (1H, broad s)

Preparation 15

To a solution of 2-benzhydryloxyacarbonylmethoxyimino-2-(2-formamidothiazol-4-yl)acetic acid (syn isomer) (7.5 g) in tetrahydrofuran (40 ml) was added trifluoroacetic anhydride (7.9 g) at -16°C for 10 minutes. To the reaction mixture was added triethylamine

(5.3 ml) at -10°C, and then the mixture was stirred

for 90 min at 0 to 5°C. To the above mixture were added ethyl acetate (100 ml) and water (100 ml), and adjusted to pH 7.5 with a saturated aqueous sodium bicarbonate. The aqueous layer was adjusted to pH 2.0

with conc. hydrochloric acid and extracted with

ethyl acetate (200 ml). The organic layer was washed with a saturated aqueous sodium chloride and dried over magnesium sulfate. The solvent was removed by evaporation under reduced pressure to give an oil. n-Hexane was added to the oil and the precipitated substance was collected by filtration to give crystalline 2-benzhydryloxyacarbonylmethoxyimino-2-[2-(2,2-trifluoroacetamido)thiazol-1-yl]acetic acid (syn isomer) (6.0 g), mp 178-180°C.

IR (Nujol): 1752, 1726 cm^{-1}

NMR δ ppm (DMSO-d_6): 4.98 (2H, s), 6.92 (1H, s), 7.32 (10H, m), 7.69 (1H, s)

(continued to the next page)

-64-

Preparation of the object compoundsExample 1

To a solution of N-tert-butoxycarbonyl-2-(3-methanesulfonamidophenyl)-D-glycine (3.3 g) and triethylamine (1.34 ml) in tetrahydrofuran (50 ml) was added dropwise a solution of ethyl chloroformate (0.91 ml) in tetrahydrofuran (10 ml) at -10 to -7°C with stirring, and the stirring was continued at the same temperature for 40 minutes to give a solution of the activated acid. This solution was added dropwise to a solution of benzhydryl 7-amino-3-methylthiomethyl-3-cephem-4-carboxylate (3.0 g) in methylene chloride (100 ml) at -30°C over a period of 5 minutes with stirring, and the stirring was continued at the same temperature for an hour. After addition of water, the reaction mixture was stirred for half an hour, followed by extracting with methylene chloride (100 ml). The extract was washed twice with 5% aqueous solution of sodium bicarbonate (50 ml) and an aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate, followed by evaporation under reduced pressure to give benzhydryl 7-[N-tert-butoxycarbonyl-2-(3-methanesulfonamidophenyl)-D-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylate (6.5 g).

I.R. (Nujol): 1780, 1710, 1680, 1152 cm^{-1}

NMR δ ppm (DMSO- d_6): 1.36 (9H, s), 1.76 (3H, s), 2.98 (3H, s), 3.3-3.8 (4H, m), 5.12 (1H, d, J=5Hz), 5.73 (1H, dd, J=5Hz, 8Hz), 6.93 (1H, s), 7.1-7.4 (14H, m), 9.93 (1H, d, J=8Hz)

834

42353D

518

-65-

Example 2

Benzhydryl 7-[N-tert-butoxycarbonyl-2-(3-methanesulfonamidophenyl)-D-glycinamido]-3-ethylthiomethyl-3-cephem-4-carboxylate (5.4 g) was obtained by reacting benzhydryl 7-amino-3-ethylthiomethyl-3-cephem-4-carboxylate (3.08 g) with N-tert-butoxycarbonyl-2-(3-methanesulfonamidophenyl)-D-glycine (3.3 g) in substantially the same manner as that of Example 1.

I.R. (Nujol): 3250, 1780, 1700, 1530, 1490, 1455 cm^{-1}

NMR δ ppm (DMSO- d_6): 0.98 (3H, t, J=7Hz), 1.38 (9H, s), 2.28 (2H, q, J=7Hz), 2.97 (3H, s), 3.4-3.8 (4H, m), 5.12 (1H, d, J=5Hz), 5.24 (1H, m), 5.73 (1H, dd, J=5Hz, 7Hz), 6.93 (1H, s), 7.0-7.7 (14H, m), 9.89 (1H, d, J=8Hz), 10.43 (1H, s)

Example 3

(1) To a suspension of tert-butyl 7-amino-3-methoxymethyl-3-cephem-4-carboxylate tosylate (13.1 g) in methylene chloride (200 ml) was added water (100 ml), followed by adjusting to pH about 6 with an aqueous solution of sodium bicarbonate. After separating out the methylene chloride layer, the remaining aqueous solution was extracted with methylene chloride (50 ml). The combined methylene chloride solution was washed with an aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate and then treated with an activated charcoal, followed by evaporation under reduced pressure to give tert-butyl 7-amino-3-methoxymethyl-3-cephem-4-carboxylate (5.7 g).

I.R. (Nujol): 3370, 1760, 1740, 1710 cm^{-1}

NMR δ ppm (DMSO- d_6): 1.5 (9H, s), 2.3 (2H, broad s), 3.24 (3H, s), 3.51 (2H, s), 4.15 (2H, s).

835

42353D

519

4.8 (1H, d, J=5Hz); 5.04 (1H, d, J=5Hz)
(2) On the other hand, to a solution of N-tert-butoxycarbonyl-2-(3-methanesulfonamidophenyl)-D-glycine (8.89 g) and triethylamine (2.61 g) in dry tetrahydrofuran (95 ml) was added dropwise a solution of ethyl chloroformate (2.8 g) in dry tetrahydrofuran (25 ml) at -10 to -7°C over a period of 10 minutes with stirring, and stirring was continued at the same temperature for 40 minutes to prepare a solution of the activated acid.

(3) To a solution of the compound (5.65 g) obtained according to Example 3-(1) in dry methylene chloride (190 ml) was added dropwise the solution of the activated acid prepared above at -30°C over a period of 10 minutes with stirring, and the stirring was continued at the same temperature for an hour.

After addition of water (100 ml), the reaction mixture was stirred for half an hour. The methylene chloride layer was separated out therefrom and the remaining aqueous solution was extracted with methylene chloride. The combined methylene chloride solution was washed twice with 5% aqueous solution of sodium bicarbonate (100 ml) and an aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate, and then treated with an activated charcoal, followed by evaporation under reduced pressure to give tert-butyl 7-[N-tert-butoxycarbonyl-2-(3-methanesulfonamidophenyl)-D-glycinamido]-3-methoxymethyl-3-cephem-4-carboxylate (17.8 g).

I.R. (Nujol): 3250, 1780, 1710, 1680 cm⁻¹

NMR 6ppm (DMSO-d₆): 1.36 (9H, s), 1.46 (9H, s), 2.97 (3H, s), 3.18 (3H, s), 3.4 (2H, broad s), 4.07 (2H, broad s), 5.03 (1H, d, J=5Hz), 5.24 (1H, d, J=8Hz), 5.73 (1H, dd, J=5Hz, 8Hz), 6.93-7.43

836

42353D

520

(4H, m), 7.5 (1H, d, J=8Hz), 9.16 (1H, d, J=8Hz), 9.73 (1H, broad s),

The compounds described in the following Examples 4 to 7 were obtained by reacting the corresponding 7-aminocephalosporanic acid derivative with 2-(3-methanesulfonamidophenyl)-D-glycine in substantially the same manner as that of Example 3.

Example 4

7-[2-(3-Methanesulfonamidophenyl)-D-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid.
I.R. (Nujol): 1758, 1687 (shoulder), 1666, 1144, 974 cm⁻¹

Example 5

7-[2-(3-Methanesulfonamidophenyl)-D-glycinamido]-3-ethylthiomethyl-3-cephem-4-carboxylic acid.
I.R. (Nujol): 3500, 3150, 1780, 1685, 1460 cm⁻¹

Example 6

7-[2-(3-Methanesulfonamidophenyl)-D-glycinamido]-3-methoxymethyl-3-cephem-4-carboxylic acid.
I.R. (Nujol): 3500, 3150, 1760, 1685 cm⁻¹

Example 7

7-[2-(3-Methanesulfonamidophenyl)-D-glycinamido]-3-allylthiomethyl-3-cephem-4-carboxylic acid tri-fluoroacetate.
I.R. (Nujol): 1760, 1680, 1600, 1140 cm⁻¹

Example 8

A mixture of benzhydryl 7-[N-tert-butoxycarbonyl-2-(3-methanesulfonamidophenyl)-D-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylate (6.5 g), anisole (5.0 ml) in trifluoroacetic acid (20 ml) was stirred at 25°C for 15 minutes. After the reaction, the reaction mixture was added dropwise to diisopropyl ether (300 ml), and the precipitated solid was collected by filtration, washed with diisopropyl ether and then dissolved in a mixture of water (50 ml) and ethyl acetate (50 ml). After the aqueous layer was

837

521

42353D

separated out, it was washed with ethyl acetate, followed by removal of ethyl acetate from the aqueous solution completely under reduced pressure.

The resultant aqueous solution was adjusted to pH 3.8 with an aqueous solution of sodium bicarbonate and subjected to column chromatography using "Diaion HP-20" (90 ml). After washing with water (180 ml), elution was carried out with 30% isopropyl alcohol, and the fractions containing the desired compound were

10

(continued to the next page)

15

20

25

30

35

838

423530

522

collected and lyophilized to give 7-[2-(3-methanesulfonamidophenyl)-D-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (4.8 g).

I.R. (Nujol): 1758, 1687 (shoulder), 1666, 1144, 974 cm^{-1}

5

NMR δ ppm ($\text{D}_2\text{O} + \text{DCI}$): 1.98 (3H, s), 3.15 (3H, s), 3.45 (2H, broad s), 3.56 (2H, broad s), 5.12 (1H, d, J=5Hz), 5.30 (1H, s), 5.70 (1H, d, J=5Hz), 7.45 (4H, s)

10

Example 9

7-[2-(3-Methanesulfonamidophenyl)-D-glycinamido]-3-ethylthiomethyl-3-cephem-4-carboxylic acid (1.6 g) was obtained by reacting benzhydrol 7-[N-tert-butoxycarbonyl-2-(3-methanesulfonamidophenyl)-D-glycinamido]-3-ethylthiomethyl-3-cephem-4-carboxylate (3.7 g) with trifluoroacetic acid (7.4 ml) in the presence of anisole (7.4 ml) in substantially the same manner as that of Example 8, mp 188°C (dec.).

15

20

I.R. (Nujol): 3500, 3150, 1780, 1685, 1460 cm^{-1}
NMR δ ppm ($\text{D}_2\text{O} + \text{DCI}$): 1.13 (3H, t, J=7Hz), 2.48 (2H, q, J=7Hz), 3.11 (3H, s), 3.3-3.8 (4H, m), 5.30 (1H, s), 5.65 (1H, d, J=5Hz), 7.43 (4H, s)

Example 10

To a solution of tert-butyl 7-[N-tert-

25

butoxycarbonyl-2-(3-methanesulfonamidophenyl)-D-glycinamido]-3-methoxymethyl-3-cephem-4-carboxylate (15 g) in anisole (11.25 ml) was added dropwise trifluoroacetic acid (33.75 ml) below 15°C over a period of 10 minutes with stirring, and the stirring was continued at 15 to 20°C for half an hour.

30

The reaction mixture was poured into diisopropyl ether (750 ml), followed by stirring at ambient temperature for 20 minutes. After the precipitated solid was collected by filtration and washed with diisopropyl

35

423530

839

523

-70-

ether, it was poured into a mixture of ethyl acetate (100 ml) and water (100 ml) and stirred for a while. The aqueous solution was separated out therefrom, and the remaining organic layer was extracted with water. The combined aqueous solution was concentrated under reduced pressure, and the concentrate was adjusted to pH about 3.8 with an aqueous solution of sodium bicarbonate, followed by subjecting to column chromatography using non-ionic adsorption resin "Diaion HP-20" (225 ml), which was washed with water (450 ml) and then eluted with 30% isopropyl alcohol.

The fractions containing the desired compound were collected and evaporated under reduced pressure, and the residue was lyophilized to give 7-[2-(3-methanesulfonamidophenyl)-D-glycinamido]-3-methoxymethyl-3-cephem-4-carboxylic acid (4.45 g).

I.R. (Nujol): 3500, 3150, 1760, 1685 cm^{-1} .

NMR δ ppm ($\text{D}_2\text{O} + \text{DCl}$): 3.13 (3H, s), 3.26 (3H, s), 3.42 (2H, q, $J=18\text{Hz}$), 4.25 (2H, s), 5.06 (1H, d, $J=5\text{Hz}$), 5.27 (1H, s), 5.73 (1H, d, $J=5\text{Hz}$), 7.42 (4H, s)

Example 11

Benzhydryl 7-amino-3-allylthiomethyl-3-cephem-4-carboxylate (3.3 g) and N-tert-butoxycarbonyl-2-(3-methanesulfonamidophenyl)-D-glycine (3.44 g) were treated in substantially the same manner as that of Example 1 to give an oily product (6.5 g).

A mixture of this oil, trifluoroacetic acid (15 ml) and anisole (15 ml) was stirred at 20°C for half an hour, followed by adding dropwise to diisopropyl ether. The precipitated solid was collected by filtration, washed with diisopropyl ether and then dissolved in a mixture of water (50 ml) and ethyl acetate (50 ml). The aqueous layer was separated out and washed with diethyl ether, followed by removal of the organic

840

42353D

524

-71-

solvent therefrom completely. The resultant aqueous solution was lyophilized to give 7-[2-(3-methanesulfonamidophenyl)-D-glycinamido]-3-allylthiomethyl-3-cephem-4-carboxylic acid trifluoroacetate (2 g).

I.R. (Nujol): 1760, 1680, 1600, 1140 cm^{-1}

NMR δ ppm ($\text{D}_2\text{O} + \text{DCl}$): 3.20 (2H, m), 3.23 (3H, s), 3.6 (7H, m), 5.2 (1H, d, $J=5\text{Hz}$), 5.45 (1H, s), 5.80 (1H, d, $J=5\text{Hz}$), 7.55 (3H, m)

Example 12

To a solution of benzhydryl 7-[4-bromoacetoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (6.0 g) in tetrahydrofuran (30 ml) was added dropwise a solution of thiourea (0.85 g) and sodium bicarbonate (0.94 g) in tetrahydrofuran (30 ml) and water (24 ml) at 25°C with stirring, and the stirring was continued at 28 to 30°C for an hour. The reaction mixture was poured into a mixture of ethyl acetate (100 ml) and water (100 ml), followed by separating out the organic layer, which was washed twice with an aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate and then evaporated to give benzhydryl 7-[2-(2-aminothiazol-4-yl)-acetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (4.0 g).

IR (Nujol): 1773, 1715, 1653 cm^{-1}

NMR δ ppm ($\text{DMSO}-d_6$): 1.85 (3H, s), 3.3-3.8 (6H, m), 5.25 (1H, d, $J=5\text{Hz}$), 5.80 (1H, dd, $J=5\text{Hz}$, 8Hz), 6.32 (1H, s), 7.00 (1H, s), 7.43 (10H, s), 8.95 (1H, d, $J=8\text{Hz}$)

Example 13

tert-Butyl 7-amino-3-methoxymethyl-3-cephem-4-carboxylate tosylate (2.0 g) was dissolved in a mixture of acetone (40 ml) and a saturated aqueous solution of sodium bicarbonate (15 ml), and thereto was added

841

42353D

525

-72-

dropwise a solution of the activated acid, which was prepared from 2-(2-formamidothiazol-4-yl)-2-methoxyiminoacetic acid (syn isomer) (1.06 g), phosphorus oxychloride (0.85 g) and N,N-dimethylformamide (0.41 g) in tetrahydrofuran (10 ml), at 0 to 5°C over a period of 10 minutes. During the addition, the pH value of the reaction mixture was maintained at pH 7.0 to 7.5 with a saturated aqueous solution of sodium carbonate. After stirring for an hour, the reaction mixture was diluted with water (50 ml), followed by extracting twice with ethyl acetate. The combined extract was washed with 5% aqueous solution of sodium bicarbonate and water, dried and then evaporated to dryness under reduced pressure to give a residue, which was triturated with diethyl ether to obtain tert-butyl 7-[2-(2-formamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate (syn isomer) (1.14 g).

IR (Nujol) : 3250, 3100, 1790, 1710, 1660 cm^{-1}
 NMR δ ppm (DMSO- d_6) : 1.49 (9H, s), 3.21 (3H, s), 3.28 (2H, broad s), 3.89 (3H, s), 4.1 (2H, s), 5.16 (1H, d, J=5Hz), 5.80 (1H, dd, J=5Hz, 8Hz), 7.36 (1H, s), 8.48 (1H, s), 9.6 (1H, d, J=8Hz), 12.66 (1H, broad s)

Example 14

tert-Butyl 7-[2-(2-formamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (syn isomer) (0.91 g) was obtained by reacting tert-butyl 7-amino-3-methylthiomethyl-3-cephem-4-carboxylate (0.90 g) with 2-(2-formamidothiazol-4-yl)-2-methoxyiminoacetic acid (syn isomer) (0.85 g) according to the similar manner to that of Example 13.

842

42353D

526

35

-73-

IR (Nujol) : 3250, 3050, 1780, 1690 cm^{-1}
 NMR δ ppm (DMSO- d_6) : 1.47 (9H, s), 1.97 (3H, s), 3.29 (2H, broad s), 3.55 (2H, ABq, J=13Hz), 3.87 (3H, s), 5.21 (1H, d, J=5Hz), 5.76 (1H, dd, J=5Hz, 8Hz), 7.38 (1H, s), 8.49 (1H, s), 9.66 (1H, d, J=8Hz), 12.56 (1H, broad s)

5

The compounds described in the following Examples 15 to 17 were obtained by reacting the 7-aminocephalosporanic acid derivatives with the corresponding acid according to the similar manner to that of Example 13.

Example 15

7-[2-(2-Aminothiazol-4-yl)acetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid
 IR (Nujol) : 1763, 1654 cm^{-1}

Example 16

7-[2-(2-Aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid hydrochloride (syn isomer)
 IR (Nujol) : 3300, 1780, 1720, 1660, 1640 cm^{-1}

20

Example 17

7-[2-(2-Aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (syn isomer)
 IR (Nujol) : 3350, 1770, 1670 cm^{-1}

Example 18

The mixture of benzhydryl 7-[2-(2-aminothiazol-4-yl)acetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (3.3 g), trifluoroacetic acid (10 ml) and anisole (10 ml) in methylene chloride (10 ml) was stirred at 10°C for an hour. After benzene (50 ml) was added to the reaction mixture, the trifluoroacetic acid therein was azeotropically removed under reduced pressure. The remained aqueous solution was poured into a mixture of ethyl acetate (100 ml) and water (100 ml), followed

35

843

42353D

527

by adjusting to pH 7.5 with sodium bicarbonate. After separating out the aqueous solution, the organic solvent was removed therefrom by evaporation completely under reduced pressure, followed by adjusting to pH 3.0 with 10% hydrochloric acid. The precipitated solid was collected by filtration and then dried to give 7-[2-(2-aminothiazol-4-yl)acetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (0.95 g).

IR (Nujol) : 1763, 1654 cm^{-1}

NMR δ ppm (DMSO- d_6) : 2.01 (3H, s), 3.48 (2H, s), 3.65 (4H, broad s), 5.17 (1H, d, J=5Hz), 5.67 (1H, dd, J=5Hz, 8Hz), 6.38 (1H, s), 9.00 (1H, d, J=8Hz)

Example 19

To a cold suspension of tert-butyl 7-[2-(2-formamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylate (syn isomer) (1.0g) in methylene chloride (20 ml) were added trifluoroacetic acid (4.4 g) and anisole (0.2 ml), and the mixture was gradually warmed at 40°C, followed by stirring at 10°C for 3 hours. After evaporation of the reaction mixture, to the residue was added ethyl acetate (20 ml), followed by extracting with an aqueous solution of sodium bicarbonate. The resultant aqueous solution was adjusted to pH 2.0 with diluted hydrochloric acid and then extracted with ethyl acetate. The extract was washed with a saturated aqueous solution of sodium bicarbonate, water and an aqueous solution of sodium chloride in turn, dried and then evaporated to dryness under reduced pressure. The residue was stirred for 30 minutes in diethyl ether, and the remained substance was collected by filtration to give 7-[2-(2-formamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid (syn isomer) (0.56 g).

844

423530

- 528

IR (Nujol) : 3250, 1780, 1660 cm^{-1}
 NMR δ ppm (DMSO- d_6) : 3.22 (3H, s), 3.55 (2H, broad s), 3.90 (3H, s), 4.19 (2H, s), 5.17 (1H, d, J=5Hz), 5.81 (1H, dd, J=5Hz, 8Hz), 7.43 (1H, s), 8.51 (1H, s), 9.67 (1H, d, J=8Hz), 12.58 (1H, broad s)

Example 20

7-[2-(2-Formamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (syn isomer) (0.64 g) was obtained by reacting tert-butyl 7-[2-(2-formamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (syn isomer) (0.9 g) with trifluoroacetic acid (5.8 g) in the presence of anisole (0.9 ml) according to the similar manner to that of Example 19.

IR (Nujol) : 3250, 1780, 1660 cm^{-1}
 NMR δ ppm (DMSO- d_6) : 1.99 (3H, s), 3.38-4.1 (4H, m), 3.93 (3H, s), 5.25 (1H, d, J=5Hz), 5.79 (1H, dd, J=5Hz, 8Hz), 7.45 (1H, s), 8.43 (1H, s), 9.65 (1H, d, J=8Hz), 12.68 (1H, broad s)

The compounds described in the following Examples 21 and 22 were obtained by reacting tert-butyl ester of the corresponding cephalosporanic acid derivatives with trifluoroacetic acid in the presence of anisole according to the similar manner to that of Example 19.

Example 21

7-[2-(2-Aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid hydrochloride (syn isomer)

IR (Nujol) : 3300, 1780, 1720, 1660, 1640 cm^{-1}

Example 22

7-[2-(2-Aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (syn isomer)

IR (Nujol) : 3350, 1770, 1670 cm^{-1} - 529

845

423530

Example 23

To a suspension of 7-[2-(2-formamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid (syn isomer) (0.52 g) in a mixture of methanol (3 ml) and tetrahydrofuran (2 ml) was added conc. hydrochloric acid (0.18 g), and the mixture was stirred at 30°C for 4 hours. After the reaction mixture was cooled and diluted with diisopropyl ether, the precipitated crystals were collected by filtration, washed with diisopropyl ether and then dried to give 7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid hydrochloride (syn isomer) (0.45 g).

IR (Nujol) : 3300, 1780, 1720, 1660, 1640 cm^{-1}
 NMR δ ppm (DMSO- d_6) : 3.26 (3H, s), 3.58 (2H, broad s), 4.0 (3H, s), 4.24 (2H, s), 5.24 (1H, d, J=5Hz), 5.82 (1H, dd, J=5Hz, 8Hz), 7.01 (1H, s), 9.87 (1H, d, J=8Hz)

Example 24

7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (syn isomer) (0.25 g) was obtained by reacting 7-[2-(2-formamidothiazol-4-yl)-2-methoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (syn isomer) (0.6 g) with conc. hydrochloric acid (0.4 g) in a mixture of methanol (3 ml) and tetrahydrofuran (1 ml) according to the similar manner to that of Example 23.

IR (Nujol) : 3350, 1770, 1670 cm^{-1}
 NMR δ ppm (DMSO- d_6) : 2.01 (3H, s), 3.2-4.1 (4H, m), 3.88 (3H, s), 5.25 (1H, d, J=5Hz), 5.78 (1H, dd, J=5Hz, 8Hz), 6.82 (1H, s), 7.24 (2H, broad s), 9.64 (1H, d, J=8Hz)

35

- 846

42353D

530

Example 25

To a solution of benzhydryl 7-amino-3-methylthiomethyl-3-cephem-4-carboxylate (10 g) and trimethylsilylacetamide (18.4 g) in methylene chloride (100 ml) was added at -15°C an activated acid, which was prepared from (2-formamidothiazol-4-yl)glyoxylic acid (6.56 g), N,N-dimethylformamide (2.92 ml) and phosphorus oxychloride (5.46 ml) in a conventional manner, and the mixture was stirred at -20 to -15°C for 15 minutes. After the reaction mixture was poured into water, it was extracted with ethyl acetate. The extract was washed with an aqueous solution of sodium bicarbonate and an aqueous solution of sodium chloride, followed by drying over anhydrous magnesium sulfate. Removal of the solvent gave a residue (13.4 g), which was chromatographed on silica gel (100 ml) using a mixture of benzene and ethyl acetate (5:1 by volume) as an eluent. The fractions containing the desired compound were collected and then evaporated to dryness to obtain benzhydryl 7-[(2-formamidothiazol-4-yl)glyoxylamido]-3-methylthiomethyl-3-cephem-4-carboxylate (7.1 g).

I.R. (Nujol) : 3300, 1780, 1700, 1656 cm^{-1}
 N.M.R. δ ppm (DMSO- d_6) : 1.83 (3H, s), 3.58 (2H, broad s), 3.67 (2H, broad s), 5.35 (1H, d, J=5Hz), 5.87 (1H, dd, J=5Hz, 8Hz), 6.90 (1H, s), 7.40 (10H, s), 8.47 (1H, s), 8.58 (1H, s), 9.88 (1H, d, J=8Hz), 12.68 (1H, broad s)

The following compounds were obtained by reacting 7-amino-3-substituted cephalosporanic acid derivatives with the corresponding acids according to the similar manner to that of Example 25.

35

- 847

42353D

- 531

0029557

Example 26

7-[(2-Aminothiazol-4-yl)glyoxylamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid.

I.R. (Nujol) : 3300, 1762, 1522 cm^{-1}

Example 27

7-[2-(2-Aminothiazol-4-yl)-DL-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid.

I.R. (Nujol) : 3300, 1755, 1686, 1600 cm^{-1} .

Example 28

To a mixture of benzhydryl 7-amino-3-methylthio-methyl-3-cephem-4-carboxylate (4.92 g) and N-tert-butoxycarbonyl-2-(2-formamidothiazol-4-yl)-DL-glycine

(4.0 g) in methylene chloride (100 ml) and tetrahydrofuran (80 ml) was added N,N'-dicyclohexylcarbodiimide (2.39 g), and the mixture was stirred at ambient temperature for an hour. After the insoluble substance was removed by filtration, the filtrate was evaporated to dryness to give a residue, which was dissolved in ethyl acetate. This solution was washed with an aqueous solution of sodium bicarbonate and an aqueous solution of sodium chloride, and then dried over anhydrous magnesium sulfate. Removal of the solvent gave benzhydryl

7-[N-tert-butoxycarbonyl-2-(2-formamidothiazol-4-yl)-DL-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylate (9.4 g).

I.R. (Nujol) : 3300, 1770, 1710, 1680, 1615 cm^{-1}

N.M.R. δ ppm (DMSO- d_6) : 1.36 (9H, s), 1.80 (3H, s), 3.58 (4H, m), 5.16 (d, J=5Hz) (1H), 5.25 (d, J=5Hz) (1H)

5.38 (1H, d, J=8Hz), 5.6-5.9 (1H, m), 6.90 (1H, s), 7.13 (1H, s), 7.33 (10H, broad s), 8.47 (1H, s), 9.08 (1H, d, J=8Hz)

Example 29

A mixture of 7-[(2-formamidothiazol-4-yl)-

glyoxylamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (3.0 g) and conc. hydrochloric acid (3 ml) in methanol (50 ml) was stirred at ambient temperature for 2 hours. The reaction mixture was adjusted to pH 5-6 with an aqueous solution of sodium bicarbonate and then concentrated under reduced pressure. The precipitated crystals in the concentrate were collected by filtration to give 7-[(2-aminothiazol-4-yl)glyoxylamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (1.1 g).

I.R. (Nujol) : 3300, 1762, 1522 cm^{-1}

N.M.R. δ ppm (DMSO- d_6) : 2.00 (3H, s), 3.65 (4H, broad s), 5.22 (1H, d, J=5Hz), 5.68

(1H, dd, J=5Hz, 8Hz), 7.37 (2H, broad s),

7.83 (1H, s), 9.73 (1H, d, J=8Hz)

Example 30

A mixture of benzhydryl 7-[N-tert-butoxycarbonyl-2-(2-formamidothiazol-4-yl)-DL-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylate (9.4 g) and conc. hydrochloric acid (4.16 ml) in methanol (100 ml) and tetrahydrofuran (25 ml) was stirred at 30 to 35°C for 5 hours. The reaction mixture was adjusted to pH 4.5 with sodium bicarbonate and then evaporated to dryness to give a

residue, which was dissolved in ethyl acetate. This solution was washed with an aqueous solution of sodium chloride, dried over anhydrous magnesium sulfate and then evaporated to obtain benzhydryl 7-[N-tert-butoxycarbonyl-2-(2-aminothiazol-4-yl)-DL-glycinamido]-3-methylthio-

methyl-3-cephem-4-carboxylate (7.4 g).

I.R. (Nujol) : 3300, 1772, 1716, 1685, 1623, 1244,

1170, 1166 cm^{-1}

N.M.R. δ ppm (DMSO- d_6) : 1.40 (9H, s), 1.80 (3H, s), 3.6 (4H, m), 5.0-5.4 (2H, m), 5.6-5.9 (2H, m), 6.93 (1H, s), 7.30 (1H, s), 7.43 (10H, broad s), 8.93 (1H, d, J=8Hz)

The following compounds were obtained by reacting 7-acylamino-3-substituted cephalosporanic acid derivatives having a formamido group with hydrochloric acid according to the similar manner to that of Example 30.

Example 31

7-[2-(2-Aminothiazol-4-yl)-DL-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid.
I.R. (Nujol) : 3300, 1755, 1686, 1600 cm^{-1}

Example 32

7-[2-(2-Aminothiazol-4-yl)-DL-glycolamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid.
I.R. (Nujol) : 3300, 1752, 1675, 1600 cm^{-1}

Example 33

Benzhydryl 7-[2-(2-formamidothiazol-4-yl)glyoxylamido]-3-methylthiomethyl-3-cephem-4-carboxylate (7.0 g) was dissolved in a solution of methylene chloride (70 ml), anisole (7 ml) and trifluoroacetic acid (14 ml), and the mixture was stirred at ambient temperature for an hour. After the solvent was removed by distillation under reduced pressure, the residue was dissolved in water, adjusted to pH 2.0 with conc. hydrochloric acid and then extracted with ethyl acetate. The extract was washed with an aqueous sodium chloride and dried over magnesium sulfate, followed by evaporation. The residue was pulverized with diisopropyl ether to obtain 7-[2-(2-formamidothiazol-4-yl)glyoxylamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (3.3 g).

I.R. (Nujol) : 3120, 1762, 1731, 1677 cm^{-1}
N.M.R. δ ppm (DMSO- d_6) : 2.00 (3H, s), 3.63 (4H, broad s), 5.24 (1H, d, J=5Hz), 5.73 (1H, dd, J=5Hz, 8Hz), 8.43 (1H, s), 8.57 (1H, s), 9.90 (1H, d, J=8Hz), 12.80 (1H, broad s)

The following compounds were obtained by reacting 7-acylamino-3-substituted cephalosporanic acid derivatives having benzhydryl ester with 2,2,2-trifluoroacetic acid in the presence of anisole according to the similar manner to that of Example 33.

Example 34

7-[(2-Aminothiazol-4-yl)glyoxylamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid.
I.R. (Nujol) : 3300, 1762, 1522 cm^{-1}

Example 35

7-[2-(2-Aminothiazol-4-yl)-DL-glycolamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid.
I.R. (Nujol) : 3300, 1752, 1675, 1600 cm^{-1}

Example 36

A mixture of benzhydryl 7-[N-tert-butoxycarbonyl-2-(2-aminothiazol-4-yl)-DL-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylate (7.0 g), anisole (7 ml) and 2,2,2-trifluoroacetic acid (21 ml) was stirred at 5°C for half an hour. To the reaction mixture was added dropwise diisopropyl ether (300 ml), and the precipitated substance was collected by filtration and then washed with diisopropyl ether, followed by dissolving in a mixture of ethyl acetate (50 ml) and water (100 ml). After the aqueous layer was separated out, the ethyl acetate in the aqueous layer was completely removed, and then the remained aqueous solution was adjusted to pH 4.2 with 5% aqueous solution of sodium bicarbonate. The resultant aqueous solution was chromatographed on nonionic adsorption resin "Diaion HP-20" (Trade Mark, made by Mitsubishi Chemical Industries Ltd.) (100 ml). After washing with water (250 ml), elution was carried out with 30% aqueous isopropyl alcohol. The fractions containing the desired compound were collected and then evaporated, followed by

lyophilization to obtain 7-[2-(2-aminothiazol-4-yl)-DL-glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (2.9 g).

I.R. (Nujol) : 3300, 1755, 1686, 1600 cm^{-1}

N.M.R. δ ppm ($\text{D}_2\text{O}-\text{DCI}$) : 2.07 (3H, s), 3.5-4.0 (4H, m), 5.27 (1H, d, J=5Hz), 5.50 (1H, s), 5.60 (d, J=5Hz), 5.72 (d, J=5Hz) (1H), 7.27 (1H, s)

Example 37

To a solution of 7-[(2-aminothiazol-4-yl)glyoxyamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (1.1 g) in methanol (80 ml) was added sodium borohydride (150 mg) at 5 to 10°C, and the mixture was stirred at the same temperature for half an hour. After the reaction mixture was adjusted to pH 5.0 with 10% hydrochloric acid, the solvent was removed by distillation. To the residue was added water (50 ml), followed by adjusting to pH 5.0 with 10% hydrochloric acid. The resultant aqueous solution was chromatographed on nonionic adsorption resin "Diaion HP-20" (50 ml). After washing with water (100 ml), elution was carried out with 30% aqueous isopropyl alcohol. The fractions containing the desired compound were collected and then evaporated to dryness, followed by lyophilization to obtain 7-[2-(2-aminothiazol-4-yl)-DL-glycolamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (0.75 g).

I.R. (Nujol) : 3300, 1752, 1675, 1600 cm^{-1}

N.M.R. δ ppm ($\text{DMSO}-d_6$) : 2.00 (3H, s), 3.48 (2H, broad s), 3.67 (2H, broad s), 4.93 (1H, s), 5.07 (1H, d, J=5Hz), 5.57 (1H, m), 6.50 (1H, s), 7.03 (2H, broad s), 8.33 (d, J=8Hz) (1H), 8.42 (d, J=8Hz) (1H)

Example 38

Vijsmeier reagent prepared from phosphorus oxychloride (1.23 ml) and N,N-dimethylformamide (1.1 ml) was suspended in dry tetrahydrofuran (40 ml). To the suspension was added 2-(2-formamidothiazol-4-yl)-2-tert-butoxycarbonylmethoxyiminoacetic acid (syn isomer) (4.0 g) under ice-cooling with stirring, and the mixture was stirred at the same temperature for 50 minutes to prepare the activated acid solution. On the other hand, benzhydryl 7-amino-3-methylthiomethyl-3-cephem-4-carboxylate (4.66 g) and trimethylsilylacetamide (8.6 g) were dissolved in methylene chloride (50 ml). To the solution was added the activated acid solution at -20°C at a time, and the mixture was stirred at the same temperature for an hour. Water (100 ml) and ethyl acetate (200 ml) were added to the resultant solution, and the organic layer was separated, washed with 5% aqueous sodium bicarbonate and an aqueous sodium chloride, and then dried over magnesium sulfate, followed by evaporation under reduced pressure to give benzhydryl 7-[2-(2-formamidothiazol-4-yl)-2-tert-butoxycarbonylmethoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (syn isomer) (8.0 g), mp 132-138°C.

IR (Nujol) : 3260, 1783, 1725, 1687 cm^{-1}

NMR δ ppm ($\text{DMSO}-d_6$, δ) : 1.47 (9H, s), 1.83 (3H, s), 3.60 (2H, broad s), 3.66 (2H, broad s), 4.98 (2H, broad s), 5.32 (1H, d, J=5Hz), 5.92 (1H, dd, J=5Hz, 8Hz), 6.95 (1H, s), 7.4 (10H, m), 7.48 (1H, s), 8.54 (1H, s), 9.63 (1H, d, J=8Hz), 12.65 (1H, broad s)

Example 39

Benzhydryl 7-[2-(2-formamidothiazol-4-yl)-2-benzhydryloxycarbonylmethoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (syn isomer)

(7.8 g) was obtained by reacting benzhydryl 7-amino-3-methylthiomethyl-3-cephem-4-carboxylate (4.0 g) with the activated acid solution prepared from 2-(2-formamidothiazol-4-yl)-2-benzhydryloxyacarbonyl-methoxyiminoacetic acid (syn isomer) (6.18 g), phosphorus oxychloride (1.42 ml) and N,N-dimethylformamide (1.21 ml), according to a similar manner to that of Example 38, mp 135-142°C.

IR (Nujol) : 3250, 1780, 1722, 1685 cm^{-1}

10 NMR δ ppm (DMSO-d₆) : 1.83 (3H, s), 3.63 (4H, m), 5.0 (2H, broad s), 5.35 (1H, d, J=5Hz), 5.98 (1H, dd, J=5Hz, 8Hz), 6.95 (1H, s), 6.98 (1H, s), 7.37 (20H, m), 7.50 (1H, s), 8.57 (1H, s), 9.82 (1H, d, J=8Hz), 12.73 (1H, broad s)

Example 40

Benzhydryl 7-[2-benzhydryloxyacarbonylmethoximino-2-(2-(2,2,2-trifluoroacetamido)thiazol-4-yl)acetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (syn isomer) (5.6 g) was obtained by reacting benzhydryl 7-amino-3-methylthiomethyl-3-cephem-4-carboxylate (3.0 g) with the activated acid solution prepared from 2-benzhydryloxyacarbonylmethoximino-2-(2-(2,2,2-trifluoroacetamido)thiazol-4-yl)acetic acid (syn isomer) (3.68 g), phosphorus oxychloride (0.89 ml) and N,N-dimethylformamide (0.75 ml), according to a similar manner to that of Example 38, mp 165-169°C.

IR (Nujol) : 3300, 1786, 1733, 1675, 1610 cm^{-1}

30 NMR δ ppm (DMSO-d₆) : 2.00 (3H, s), 3.56 (4H, m), 4.84 (2H, s), 5.26 (1H, d, J=5Hz), 5.86 (1H, dd, J=5Hz, 8Hz), 6.84 (1H, s), 6.88 (1H, s), 7.3 (20H, m), 9.62 (1H, d, J=8Hz)

Example 41

To a suspension of benzhydryl 7-[2-(2-

854

42353D

538

855

42353D

539

formamidothiazol-4-yl)-2-tert-butoxycarbonylmethoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (syn isomer) (8.0 g) in methanol (160 ml) was added conc. hydrochloric acid (5.6 ml), and the mixture was stirred at 35°C for an hour.

The resultant solution was adjusted to pH 5.0 with a saturated aqueous sodium bicarbonate. After distilling methanol under reduced pressure, the residue was dissolved in water (100 ml) and ethyl acetate (200 ml).

10 The ethyl acetate layer was washed with a saturated aqueous sodium chloride and dried over magnesium sulfate. The solvent was removed by filtration to give

benzhydryl 7-[2-(2-aminothiazol-4-yl)-2-tert-

butoxycarbonylmethoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (syn isomer) (7.0 g), mp 140-145°C.

IR (Nujol) : 3250, 1780, 1723, 1680 cm^{-1}

20 NMR δ ppm (DMSO-d₆) : 1.43 (9H, s), 1.83 (3H, s), 3.63 (4H, m), 4.6 (2H, broad s), 5.29 (1H, d, J=5Hz), 5.87 (1H, dd, J=5Hz, 8Hz), 6.83 (1H, s), 6.95 (1H, s), 7.4 (10H, m), 9.52 (1H, d, J=8Hz)

Example 42

Benzhydryl 7-[2-(2-aminothiazol-4-yl)-2-

benzhydryloxyacarbonylmethoxyiminoacetamido]-3-

25 methylthiomethyl-3-cephem-4-carboxylate (syn isomer) (7.0 g), mp 148-155°C, was obtained by reacting

benzhydryl 7-[2-(2-formamidothiazol-4-yl)-2-

benzhydryloxyacarbonylmethoxyiminoacetamido]-3-

methylthiomethyl-3-cephem-4-carboxylate (syn isomer)

(7.5 g) with conc. hydrochloric acid (3.9 ml) accord-

ing to a similar manner to that of Example 41.

NMR δ ppm (DMSO-d₆) : 1.83 (3H, s), 3.60 (4H, m),

4.90 (2H, broad s), 5.30 (1H, d, J=5Hz),

5.87 (1H, dd, J=5Hz, 8Hz), 6.86 (1H, s),

6.93 (1H, s), 6.97 (1H, s), 7.4 (20H, m),

9.67 (1H, d, J=8Hz).

854

42353D

538

855

42353D

539

0020557

-86-

Example 43

A solution of 7-[2-carboxymethoxyimino-2-{2-(2,2-trifluoroacetamido)thiazol-4-yl}acetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (syn isomer) (4.1 g) and sodium acetate (9.56 g) in water (41 ml) was stirred for 19.5 hours at ambient temperature. The resultant solution was adjusted to pH 2.0 with conc. hydrochloric acid. The precipitates were collected by filtration and dried over phosphorus pentoxide to give 7-[2-carboxymethoxyimino-2-(2-aminothiazol-4-yl)acetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (syn isomer) (1.8 g), mp 173-176°C (dec.).

IR (Nujol) : 3370, 1772, 1670 (broad) cm^{-1}

NMR δ ppm (DMSO- d_6) : 2.00 (3H, s), 3.57 (4H, m),

4.60 (2H, broad s), 5.17 (1H, d, J=5Hz), 5.73

(1H, dd, J=5Hz, 8Hz), 6.77 (1H, s), 9.45 (1H, d, J=8Hz)

Example 44

To a solution of benzhydryl 7-[2-benzhydryloxy-carboxymethoxyimino-2-(2-(2,2-trifluoroacetamido)-thiazol-4-yl)acetamido]-3-methylthiomethyl-3-cephem-4-carboxylate (syn isomer) (5.6 g) and anisole (5.6 ml) in methylene chloride (11.2 ml) was added trifluoroacetic acid (11.2 ml) at 10°C. The mixture was stirred for 1.5 hours at ambient temperature and then poured into a mixture of diisopropyl ether (400 ml) and petroleum ether (100 ml). The precipitates were collected by filtration and washed with petroleum ether to give 7-[2-carboxymethoxyimino-2-(2-(2,2-trifluoroacetamido)-thiazol-4-yl)acetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid (syn isomer) (4.2 g), mp 183-186°C (dec.).

IR (Nujol) : 3300, 1778, 1723, 1660 cm^{-1}

NMR δ ppm (DMSO- d_6) : 2.00 (3H, s), 3.62 (4H, m),

4.70 (2H, s), 5.24 (1H, d, J=5Hz),

5.80 (1H, dd, J=5Hz, 8Hz), 7.60 (1H, s),


9.68 (1H, d, J=8Hz)

0020557


-87-

What we claim is:

1. A compound of the formula:



In which R^1 is a group of the formula:



wherein R^4 is lower alkyl and

R^5 is amino or a protected amino group,

R^2 is lower alkoxyethyl, lower alkyl-

thiomethyl or lower alkenylthiomethyl,

R^3 is carboxy or a protected carboxy group,

and


A is lower alkylene which may have a substituent selected from the groups consisting of amino, a protected amino group, hydroxy, oxo and a group of the formula:

-N-OR^6 , wherein R^6 is hydrogen, lower alkenyl, lower alkynyl, lower alkyl, or lower alkyl substituted by one or more substituent(s) selected from carboxy, a protected carboxy group, amino, a protected amino group and a heterocyclic group, and

a pharmaceutically acceptable salt thereof.

2. A compound of claim 1, in which

the formula: R^1 -A- is a group of the formula:



In which R^1 is a group of the formula:

858


42353D

540

857

42353D

541



, wherein R⁵ is amino or acylamino,

R⁶ is lower alkyl, carboxy(lower)alkyl or esterified carboxy(lower)alkyl, and R³ is carboxy or an esterified carboxy group.

3. A compound of claim 2, which is syn isomer.

4. A compound of claim 3, in which R⁵ is amino.

5. A compound of claim 4, in which

R² is lower alkoxyethyl or lower alkylthiomethyl,

R³ is carboxy, and

R⁶ is lower alkyl or carboxy(lower)alkyl.

6. A compound of claim 5, which is selected from the group consisting of:


7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid,

7-[2-(2-aminothiazol-4-yl)-2-methoxyiminoacetamido]-3-methoxymethyl-3-cephem-4-carboxylic acid or its hydrochloride, and

7-[2-(2-aminothiazol-4-yl)-2-carboxymethoxyiminoacetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid.

7. A compound of claim 1, in which

R¹ is a group of the formula:



wherein R⁵ is amino or acylamino,

A is methylene, aminomethylene, acylaminomethylene,

hydroxymethylene or carbonyl, and

R³ is carboxy or an esterified carboxy group.

8. A compound of claim 7, in which

R⁵ is amino.

9. A compound of claim 8, in which

R² is lower alkylthiomethyl, and

R³ is carboxy.

10. A compound of claim 9, which is selected from the group consisting of:

7-[2-(2-aminothiazol-4-yl)acetamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid,


7-[2-(2-aminothiazol-4-yl)glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid,

7-[2-(2-aminothiazol-4-yl)glycolamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid and

7-[2-(2-aminothiazol-4-yl)glyoxylamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid.

11. A compound of claim 1, in which

R¹ is a group of the formula:



wherein R⁴ is as defined


in claim 1,

A is aminomethylene or acylaminomethylene, and

R³ is carboxy or an esterified carboxy group.

12. A compound of claim 11, in which

R¹ is a group of the formula:



wherein R⁴ is as defined above.

13. A compound of claim 12, in which

R³ is carboxy.

14. A compound of claim 13, which is selected from the group consisting of:

7-[2-(3-methanesulfonamidophenyl)glycinamido]-3-methylthiomethyl-3-cephem-4-carboxylic acid,


7-[2-(3-methanesulfonamidophenyl)glycinamido]-3-ethylthiomethyl-3-cephem-4-carboxylic acid,

7-[2-(3-methanesulfonamidophenyl)glycinamido]-3-methoxymethyl-3-cephem-4-carboxylic acid and

7-[2-(3-methanesulfonamidophenyl)glycinamido]-3-allylthiomethyl-3-cephem-4-carboxylic acid or its trifluoroacetate.

15. A process for preparing a compound of the formula:

-90-



in which R^1 is a group of the formula:



wherein R^4 is lower alkyl and

R^5 is amino or a protected amino group,

R^2 is lower alkoxyethyl, lower alkyl-

thiomethyl or lower alkenylthiomethyl,

R^3 is carboxy or a protected carboxy group

and

A is lower alkylene which may have a

substituent selected from the groups

consisting of amino, a protected amino

group, hydroxy, oxo and a group of the

formula:

$-N-OR^6$, wherein R^6 is hydrogen, lower

alkenyl, lower alkynyl, lower alkyl,

or lower alkyl substituted by one or

more substituent(s) selected from

carboxy, a protected carboxy group,


amino, a protected amino group and a

heterocyclic group,

and pharmaceutically acceptable salt thereof,

which comprises

(1) reacting a compound of the formula:




423530

860


544

-91-

in which R^2 and R^3 are each as defined above,
or its reactive derivative at the amino group or
a salt thereof with a compound of the formula:




in which R^1 and A are each as defined above,
or its reactive derivative at the carboxy group
or a salt thereof to give a compound of the formula:



in which R^1 , R^2 , R^3 and A are each as defined
above,

or a salt thereof; or


(2) subjecting a compound of the formula:



in which A^1 is lower alkylene having a protected
amino group; and

R^1 , R^2 and R^3 are each as defined above,
or a salt thereof to removal reaction of the

amino-protective group to give a compound of the
formula:



in which A^2 is lower alkylene having an amino
group, and


R^1 , R^2 and R^3 are each as defined above,
or a salt thereof; or

423530


861

545

(3) subjecting a compound of the formula:




in which R_a^1 is a group of the formula:




wherein R^8 is a protected amino group, and

R^2 , R^3 and A are each as defined above, or a salt thereof to removal reaction of the amino-protective group to give a compound of the formula:




in which R_b^1 is a group of the formula:




R^2 , R^3 and A are each as defined above, or a salt thereof; or

(4) subjecting a compound of the formula:




in which R^1 is a protected carboxy group, and R^1 , R^2 and A are each as defined above, or a salt thereof to removal reaction of the carboxy-protective group to give a compound of the formula:




in which R^1 , R^2 and A are each as defined above, or a salt thereof; or

(5) introducing a carboxy-protective group into a compound of the formula:




in which R^1 , R^2 and A are each as defined above, or a salt thereof to give a compound of the formula:



in which R^1 , R^2 , R^3 and A are each as defined above, or a salt thereof; or

(6) reducing a compound of the formula:




in which A^3 is lower alkylene having an oxo group, and


R^1 , R^2 and R^3 are each as defined

above, or a salt thereof to give a compound of the formula:


-94-




in which A^4 is lower alkylene having a hydroxy group, and
 R^1 , R^2 and R^3 are each as defined above,
 or a salt thereof; or
 (7) reacting a compound of the formula:



in which A^5 is lower alkylene which may have a group of the formula: $-N-OR^6$,
 wherein R^6 is as defined above,
 X^1 is halogen, and
 R^2 and R^3 are each as defined above,
 or a salt thereof with a compound of the formula:




in which R^5 is as defined above,
 to give a compound of the formula:



in which R^2 , R^3 , R^5 and A^5 are each as defined above,
 or a salt thereof; or
 (8) subjecting a compound of the formula:

-95-




in which A^6 is lower alkylene having a group of the formula:



wherein R^6 is lower alkyl substituted by a protected carboxy group, and
 R^1 , R^2 and R^3 are each as defined above,

or a salt thereof to removal reaction of the carboxy-protective group to give a compound of the formula:




in which A^7 is lower alkylene having a group of the formula:



wherein R^6_b is lower alkyl substituted by carboxy, and
 R^1 , R^2 and R^3 are each as defined above,


or a salt thereof; or
 (9) subjecting a compound of the formula:



in which R^3 is lower alkoxycarbonyl substituted by protected amino and protected carboxy groups, and

R^1 , R^2 and A are each as defined above,


or a salt thereof to removal reaction of the amino- and carboxy-protective groups to give a compound of the formula:



in which R^3 is lower alkoxy-carbonyl substituted by amino and carboxy, and

R^1 , R^2 and A are each as defined above, or a salt thereof, or


(10) subjecting a compound of the formula:



in which A^8 is lower alkylene having a group of the formula: $-N-OR^6$, wherein R^6 is lower

alkoxy-carbonyl(lower)alkyl substituted by protected amino and protected carboxy groups or lower alkyl substituted by protected amino and protected carboxy groups, and


R^1 , R^2 and R^3 are each as defined above, or a salt thereof to removal reaction of the amino- and carboxy-protective groups to give a compound of the formula:



in which A^9 is lower alkylene having a group of the formula: $-N-OR^6$, wherein


R^6 is lower alkoxy-carbonyl(lower)-alkyl substituted by amino and carboxy or lower alkyl substituted by amino and carboxy, and R^1 , R^2 and R^3 are each as defined above, or a salt thereof; or

(11) introducing a carboxy-protective group into a compound of the formula:



in which R^1 , R^2 , R^3 and A^7 are each as defined above,

or a salt thereof to give a compound of the formula:




in which R^1 , R^2 , R^3 and A^6 are each as defined above,

or a salt thereof; or


(12) reacting a compound of the formula:

-98-




in which A¹⁰ is lower alkylene having a group of the formula: -N-OR_e^6 , wherein R_e⁶ is lower alkyl substituted by a group of the formula: $\begin{array}{c} \diagup \text{N} \diagdown \\ \diagdown \text{N} \diagup \end{array}$, and R¹, R² and R³ are each as defined above,


or a salt thereof with a compound of the formula:



in which R⁷ is lower alkyl, to give a compound of the formula:



in which A¹¹ is lower alkylene having a group of the formula: -N-OR_f^6 , wherein R_f⁶ is lower alkyl substituted by a group of the formula:



wherein R⁷ is as defined above, and R¹, R² and R³ are each as defined above,


808

42353D


552

-99-

or a salt thereof; or
(13) reacting a compound of the formula:



in which R¹, R² and A¹¹ are each as defined above, or a salt thereof with a base to give a compound of the formula:



in which A¹² is lower alkylene having a group of the formula: -N-OR_g^6 , wherein R_g⁶ is lower alkyl substituted by a cation of the formula: $\begin{array}{c} \diagup \text{N} \diagdown \\ \diagdown \text{N} \diagup \end{array}$, wherein R⁷ is as defined above, and R¹ and R² are each as defined above,

or a salt thereof.

16. A pharmaceutical composition comprising, as active ingredients, the compounds of the claim 1, in admixture with pharmaceutically acceptable carriers.


17. A method for treating an infectious disease caused by pathogenic microorganisms, which comprises administering a compound of the claim 1 to infected human being and animals.

18. A compound of the formula:

869

42353D

553



5

in which R^A is amino or a protected amino group,

$R^{b'}$ is lower alkenyl,


X is -S- or -SO-, and

R^3 is carboxy or a protected carboxy group,

and a salt thereof.

10

19: A process for preparing a compound of the formula:



15

in which R^A is amino or a protected amino group,

$R^{b'}$ is lower alkenyl,


X is -S- or -SO-, and

R^3 is carboxy or a protected carboxy group, and a salt thereof,

20

which comprises

(1) reacting a compound of the formula:



25

in which R^A is a protected amino group,

Y' is a conventional group which is

capable to be replaced by the

residue (-SR^{b'}) of the compound of

the formula: HS-R^{b'}, in which $R^{b'}$

is as defined above, and R^3 is as

defined above,

35

or a salt thereof with a compound of the formula:




5

in which $R^{b'}$ is as defined above,

or its reactive derivative at the mercapto group


to give a compound of the formula:



10


in which R^A , $R^{b'}$ and R^3 are each as defined above, or a salt thereof; or

15 (2) reducing a compound of the formula:



20

in which R^A , $R^{b'}$ and R^3 are each as defined above, or a salt thereof to give a compound of the formula:




25

in which R^A , $R^{b'}$ and R^3 are each as defined above,

or a salt thereof; or

(3) subjecting a compound of the formula:

35




5

in which R^a, R^b, and R³ are each as defined above,

or a salt thereof to removal reaction of the amino-protective group to give a compound of the formula:

10



15

in which R^b and R³ are each as defined above, or a salt thereof.

20

25

30

42353D

35

872

556